

DIRECTIONS FOR USE CONTINUED FROM BACK PANEL

azaleas, camellias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants

Note: Mosquitos coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions

Attention (Lawn and Ornamental Plant Pests): Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water.

Area Control of Ticks and Chiggers: Use EAGLE 4E CHLORPYRIFOS to control ticks and chiggers infesting non-cropland areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply EAGLE 4E CHLORPYRIFOS in water at the rate of 1/2 pint per acre (equivalent to 1/2 fl. oz. per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage usually 40 to 100 gallons per acre.

Attention (Area Control of Ticks and Chiggers): Do not allow public use of treated areas during application or until spray has dried


DOSAGES and DILUTIONS for USE on ORNAMENTAL PLANTS

Pests Controlled	Amount of EAGLE 4E CHLORPYRIFOS in Water to Make		
	3 gallons	10 gallons	100 gallons
Aphids	0.25 fl. oz.	0.8 fl. oz.	0.5 pint
Bagworms			(Equivalent to 4 ounces of chlorpyrifos per 100 gallons)
Eastern tent caterpillar			
Fall webworm			
Grasshoppers, Mites,			
Orangestriped oakworm			
Spittlebugs, Whiteflies			
Yellownecked caterpillar			
Cutworms	0.5 fl. oz.	1.6 fl. oz.	1 pint
Leafhoppers			(Equivalent to 8 ounces of chlorpyrifos per 100 gallons)
Mealybugs			
Mimosa webworm			
Redhumped caterpillar			
Thrips (exposed)			
Ash borer	1.0 fl. oz.	3.2 fl. oz.	1 quart
Lilac borer			(Equivalent to 1 pound of chlorpyrifos per 100 gallons)
Scale crawlers			
Oak lecanium			
Pine needle, Tea,			
White Peach			

DOSAGES and DIRECTIONS for USE on TURF

Pest	Dosage EAGLE 4E CHLORPYRIFOS per 1000 sq. ft.	Remarks
Ants	1/4 fl. oz.	Spray when pests first appear, retreat when needed. For sod webworms delay watering or mowing the treated area for 12 to 24 hours after treatment.
Armyworms		
Brown dog tick		
Chiggers		
Chinch bugs		
Clover mite		
Crickets		
Cutworms		
Earwigs		
Fleas		
Grasshoppers		
Sod webworms		
Turfgrass (Hyperodes) weevil	1 1/2 fl. oz.	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service Specialist for your area.
White grubs (European chafer and Japanese beetle larvae only)	1 1/2 to 3 fl. oz.	For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.

1 1/4 fl. oz. per 1000 sq. ft. = 3 3/4 fl. oz. per 5000 sq. ft. 1/2 pint per 10,700 sq. ft. = 1 quart per acre



4E CHLORPYRIFOS INSECTICIDE

ACTIVE INGREDIENTS:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate)	44.8%
Xylene	22.0%
INERT INGREDIENTS	33.2%
TOTAL	100.0%

This Product Contains 4 Pounds of Chlorpyrifos per Gallon

KEEP OUT OF REACH OF CHILDREN

WARNING

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. COMBUSTIBLE. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

To Be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts, and Other Trained Personnel Responsible for Insect Control Programs

EPA Est. No. 38621-CA

EPA Reg. No. 38621-6

FOSTER MANUFACTURING, INC.
1577 FIRST STREET COACHELLA, CA 92236 (714) 398-6154

NET CONTENTS

BEST DOCUMENT AVAILABLE 3

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

WARNING

KEEP OUT OF REACH OF CHILDREN. MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. MAY BE INJURIOUS TO EYES AND SKIN.

Do not take internally. Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing before reuse. Avoid breathing vapors and spray mist. Handle concentrate in ventilated area. Avoid spraying food crops.

STATEMENT OF PRACTICAL TREATMENT

CAUTION: Never give fluids or induce vomiting if patient is unconscious or having convulsions.

If Swallowed — Drink 1 or 2 glasses of water and induce vomiting by pumping back of throat with finger or blunt object. Do not induce vomiting by anything by mouth to an unconscious person. Get medical attention.

If On Skin — Wash skin with soap and water.

If In Eyes — Flush contaminated eyes with flowing water for at least 15 minutes and get medical attention.

Note To Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

EAGLE 4E CHLORPYRIFOS is toxic to fish, birds, and other wildlife. Keep away from lakes, streams, ponds, tidal marshes and estuaries. Shrimp and oysters may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of mixtures. Apply this product only as specified on this label.

PHYSICAL-CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

Do not use in violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow children or pets to contact treated areas or surfaces until spray has dried.

STORAGE AND DISPOSAL

STORAGE: Keep away from food, feedstuffs, and domestic water supplies. Keep container closed. Do not store near heat or open flame.

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Dumping is prohibited.

Do not reuse spray mixture or residue that cannot be used or chemically processed should be disposed of according to procedures approved by Federal, State or local disposal authorities.

Do not reuse empty container. Dispose of in an approved landfill or bury in a secure place.

MOSQUITOES: Apply EAGLE 4E CHLORPYRIFOS to mosquito infested, trap areas according to the conditions and rates shown in the following table. Mix EAGLE 4E CHLORPYRIFOS with either water or oil, such as kerosene or No. 2 fuel oil, using sufficient spray volume to give uniform coverage of area treated. Apply the spray using suitable application equipment such as hand and power ground sprayers, mist applicators, and aerial equipment.

Labeling (Mosquito Control): Read complete Environmental and use Precautions and consult your State Fish and Game Department before using this product for mosquito control.

DIRECTIONS FOR USE CONTINUED TO BACK PANEL

FOR USE TO PROTECT SOUTHERN PINE SEEDLINGS FROM DAMAGE BY PALES AND PITCH-EATING WEEVILS:

EAGLE 4E CHLORPYRIFOS insecticide is recommended for treatment of southern pine seedlings used in forest regeneration projects to reduce injury caused by damage from pales and pitch-eating weevils. Treat the seedlings immediately after transplanting by applying EAGLE 4E CHLORPYRIFOS as a foliar spray containing 2% chlorpyrifos. Prepare the spray by thoroughly mixing 5 1/2 fluid ounces of EAGLE 4E CHLORPYRIFOS per gallon of water and apply using hand or power operated spray equipment. Treat seedling with enough spray to thoroughly wet the foliage and stems to prevent runoff but do not use more than 6 gallons of spray per acre.

SEI Seller warrants that the product conforms to its chemical description and is suitably fit for the purposes stated on the label when used in accordance with the terms under normal conditions of use, but neither this warranty nor any other implied or MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and seller assumes the risk of any such use.

Patent No. 3,244,586

DIRECTIONS FOR USE CONTINUED FROM SIDE PANEL

CONDITIONS and RATES to USE for MOSQUITO CONTROL

Condition of Area to be Treated	Amount to Use Per Acre		Number Acres 1 gallon of EAGLE 4E CHLORPYRIFOS Will Treat
	EAGLE 4E CHLORPYRIFOS (fluid ounces)	Chlorpyrifos (pounds)	
Larval Control			
None to medium vegetative cover	0.4 to 0.8	0.0125 to 0.025	320 to 160
Medium to heavy vegetative cover	0.8 to 1.6+	0.025 to 0.05+	160 to 80
Adult Control			
Light to medium vegetative cover	0.8	0.025	160
Medium to heavy vegetative cover	1.6+	0.05+	80

*The higher dosage will also give faster knockdown and longer residual control where there is little or no vegetative cover.

PEST INDOORS: General Information: Use EAGLE 4E CHLORPYRIFOS insecticide to control the pests indicated in the areas listed below by application as a 0.25% or 0.5% chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute EAGLE 4E chlorpyrifos with water or with a suitable base oil* as shown in the following SPRAY DILUTION CHART. Repeat the treatment as needed. *Use only deodorized kerosene for oil-based sprays to be applied in food handling equipment.

SPRAY DILUTION CHART

Fluid Ounces† of EAGLE 4E CHLORPYRIFOS to Use		
Gallons of Spray Mixture Desired	0.25% spray	0.50% spray
Water-base Spray:		
1	2 1/2	1 1/4
3	7 1/2	4
5	12 1/2	6 1/4
10	25	12 1/2
Oil-base spray††:		
1	1 1/2	1
3	4 1/2	3
5	7 1/2	5
10	15	10 1/2

†Note: 1 fluid ounce = 2 tablespoonsful = 6 teaspoonfuls

††Calculated on basis of deodorized kerosene

Note: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture containing EAGLE 4E CHLORPYRIFOS insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays, use pyrethrins at 0.25% in water-base and 0.05% in oil-base sprays. Consult the labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use EAGLE 4E CHLORPYRIFOS in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

Application Within Food Handling Establishments (Places Other Than Private Residences in Which Food is Held, Processed, Prepared or Served), Including, But Not Limited to, Restaurants, Grocery Stores, Bakeries, Bottling Plants, Canneries, and Grain Mills: Use as a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets; along and behind baseboards; beneath and behind stoves; refrigeration units and similar food preparation areas and equipment; floor drains; and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pin stream of

insecticide should be used. Repeat treatment as needed but not more often than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back): EAGLE 4E CHLORPYRIFOS may be applied after 2 days from last treatment but emergency use should be limited to once per month.

Application of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

Application Within Residential Buildings Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional, and Commercial Buildings Including Hospitals, Stores, Manufacturing Plants and Warehouses: For Spot Treatment Only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets; around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

For control of carpet beetles, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

For control of brown dog ticks, thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters; nearby cracks and crevices, along and behind baseboards; window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas use only oil-base sprays. Prior to application de-energize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

PESTS OUT-OF-DOORS: General Information: Use EAGLE 4E CHLORPYRIFOS for control of certain pests on outside surfaces of buildings and on lawns and ornamental plants, for area control of ticks and chiggers and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

Pests on Outside Surfaces of Buildings: Use EAGLE 4E CHLORPYRIFOS at the rate of 1 1/2 fluid ounces per one gallon of water (0.5% chlorpyrifos emulsion) to control ants, clover mites, cockroaches, crickets, earwigs, fleas, flies, millipedes, mosquitos, sowbugs, spiders and ticks by application as a residual spray or with a paint brush to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

Note: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 1/2 pint of EAGLE 4E CHLORPYRIFOS per 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1000 sq. ft. to thoroughly and uniformly wet the band area.

Pests of Ornamentals: Use EAGLE 4E CHLORPYRIFOS to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with the pests listed in the following table. Dilute EAGLE 4E CHLORPYRIFOS with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7-to 10-day intervals, if needed. Note—Do not use on

DIRECTIONS FOR USE CONTINUED TO SIDE PANEL

FOSTER MANUFACTURING, INC.

1577 FIRST STREET, COACHELLA, CA 92236 (714) 398-6154

TEST DOCUMENT AVAILABLE