

38621-9

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS) WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes. Do not get on clothing. Wash thoroughly after handling. Do not use in human habitation. Do not use in areas where children or pets go. Do not get on sprayed grass until spray has completely dried.

When used in dwellings, care should be taken to avoid deposits which could be hazardous to children. Do not allow children to play in treated areas until surfaces are dry. Do not contaminate food that contains or is coming into contact with.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not use in streams and ponds. Do not apply where runoff is likely to occur. Do not apply to water or lawns. Do not use in areas where birds feed. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to it. The threat of loss of bees remains in the treated area. Do not use when bees are active. Do not use on crops or plants which are being pollinated. Do not use on plants which are being pollinated. Do not use on plants which are being pollinated.

PHYSICAL OR CHEMICAL HAZARDS

Do not use on or near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not permit children or pets to go onto sprayed grass until spray has completely dried. Do not allow children in treated areas until surfaces are dry.

STORAGE AND DISPOSAL

Storage — Do not use open flame or heat near treated area. Do not use open flame or heat near treated area. Do not use open flame or heat near treated area.

Pesticide Disposal — Pesticide spray mixture or residue that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State, or local disposal authorities.

Container Disposal — Do not reuse empty container. Do not reuse empty container. Do not reuse empty container.

DIRECTIONS FOR USE CONTINUED ON BACK PANEL

EAGLE DIAZINON® 4E INSECTICIDE

For control of cockroaches, certain other household pests, and turf and ornamental pests.

ACTIVE INGREDIENTS:

O, O-diethyl O (2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 47.5%
Xylene 26.2%
INERT INGREDIENTS 26.3%

TOTAL 100.0%

THIS PRODUCT CONTAINS 4 LBS. DIAZINON® PER GALLON
FOR USE BY PEST CONTROL OPERATORS ONLY

KEEP OUT OF REACH OF CHILDREN
WARNING

STATEMENT OF PRACTICAL TREATMENT

If Swallowed — Call a physician immediately.
If Inhaled — Remove victim to fresh air and apply respiration if needed.
If on Skin — Wash promptly with soap and water. Rinse thoroughly.
If in Eyes — Rinse eyes for at least 15 minutes with water and call a physician immediately.
Note to Physician: 2-PAM is antidotal and may be administered in conjunction with atropine sulfate.

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

DIAZINON® Trademark of CIBA GEIGY

MANUFACTURED BY

FOSTER MANUFACTURING, INC.

1577 FIRST STREET, COACHELLA, CALIFORNIA 92236

(714) 333-6711

EPA Reg. No. 38621-9 EPA Est. No. 11017-CA-01

NET CONTENTS

DIRECTIONS FOR USE, CONTINUED

EAGLE DIAZINON 4E is a systemic insecticide of Diazinon type. It is used by pest control operators after dilution with water. Use is specified on this label.

EAGLE DIAZINON 4E is not intended for use by homeowners. Homeowners should use EAGLE DIAZINON 4E.

Note: The solvent used in this concentrate may stain certain materials such as wall paper and fabrics. Care should also be taken to avoid staining.

HOUSEHOLD USE

Do not use in food preparation areas, restaurants, or other areas where food is prepared or processed. Do not use in exposed areas.

SPRAY DILUTION

Dilute EAGLE DIAZINON 4E with water.

EAGLE DIAZINON 4E spray and application

Gals. of Finished Spray	Amount EAGLE DIAZINON 4E (4 lbs. per gallon)
1	1 1/4 fl. oz.
2	2 1/2 fl. oz.
3	3 3/4 fl. oz.
10	11 3/4 fl. oz.
25	29 3/4 fl. oz.

12 1/2 tablespoons = 1 1/4 fl. oz.

Cockroaches (Waterbug), Ants, and other household pests. Apply with a pump sprayer to baseboards, cracks and crevices, under cabinets, refrigerator, etc. These insects feed on these insects.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

For heavy cockroach infestation, a strength of 1 1/2% EAGLE DIAZINON 4E is recommended.

BEST DOCUMENT AVAILABLE



E
NON® 4E
CTICIDE

cockroaches, certain other
and turf and ornamental peats.

INGREDIENTS:

propyl-8-methyl-4-	
prothioate	47.5%
.....	28.5%
16:	24.0%

TOTAL 100.0%

NET CONTAINS 4 LBS.
GAL. PER GALLON
CONTROL OPERATORS ONLY

TEACH OF CHILDREN
TRAINING

7. PRACTICAL TREATMENT

physician immediately.
 victim to fresh air and apply
 gently with soap and water. Rinse

as for at least 15 minutes with
20 or more daily.
15 is total and may be ad-
ed with atropine sulfate.

FOR ADDITIONAL
STATEMENTS

Mark of 25A-GE.GY

MANUFACTURED BY:
MARTIN, INC.

100-21711, CALIFORNIA #223
702333

EPA Reg. No. 11017-CAD-01

475

DIRECTIONS FOR USE, CONTINUED

EAGLE DIAZINON 4E is an emulsifiable solution containing 4 lbs. of Diazinon per gallon. It is intended for use by pest control operators after dilution in water for the control of those insects specified on this label.

EAGLE DIAZINON 4E is not intended for use in preparing oil
base sprays. Pest control operators who prefer oil base sprays
should use EAGLE DIAZINON 4S.

Note: The solvent used in this formulation is a petroleum distillate which may stain certain plastic, rubber, and asphalt materials such as tiles and floor coverings. Do not treat such materials. Care should also be taken to avoid spotting of wallpaper and fabrics.

HOUSEHOLD INSECTS

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas where food is exposed.

SPRAY DILUTION CHART

Dilute EAGLE DIAZINON 4E with water to make a 1%, or 1% EAGLE DIAZINON spray, and apply as indicated below.

Gals of First-cod Spray	Amount EAGLE DAZINON 4E to use for	
	1/2" Spray	1" Spray
1	1 1/2 fl oz	2 1/2 fl oz
2	2 1/2 fl oz	5 fl oz
3	3 1/2 fl oz	7 1/2 fl oz
10	12 1/2 fl oz	25 fl oz (1 1/2 pints)
25	32 fl oz (2 pints)	64 fl oz (4 pints)

*25. $\log_{10} 1000 = 3$ or

Cochlosiphum (Waldbog's Ams. Silverfish) - Make spot applications of 1% EAGLE DIAZINON as a curative spray or apply with a paint brush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves, and similar areas where these insects hide.

For heavy cockroach infestations, make the first application at a strength of 1% EAGLE DIAZINON, followed by monthly applications of 1/4% EAGLE DIAZINON.

Carpenter Beetles — Apply a 4% EAGLE DIAZINON spray in spot applications along baseboards and edges of carpeting, under carpeting and rugs and furniture, in closets and on skirting, and wherever these insects are seen or suspected. **Spiders and Scorpions** — Apply a 4% EAGLE DIAZINON spray around windows, doors, and along baseboards. Make spot applications to other areas over which these pests may crawl.

Partly $\frac{1}{2}$ Exposed Stages (larva, hooked grub beetle, Flour Beetle, *Tenebrio*, Cigarette Beetle, Drugstore Beetle, Indian Meal Moth) — Apply a 1% spray to shelves and cupboards. If the flour contains suspected of being infested it is stored in airtight containers and treated before spraying. Do not use the food packages until spray has dried. Control when necessary by spraying. This treatment is supplemental to good housekeeping and will have no effect on insects breeding.

Within food packages. Used food packages should be destroyed.

Brown dog ticks in buildings — Make thorough application of a 3% EAGLE DACTHION spray to infested areas around vestiboles, windows and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and rug coverings. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

ORNAMENTAL INSECTS

For use on Azaleas, Azalea Bush, Boragid, Camellia, Car
nation, Chrysanthemum, Douglas fir, Elm, Gladiolus, Hawthorn,
Holly, Juniper, Lilac, Locust, Magnolia, Oak, Pine, Plum, Pteris,
Rhododendron, Rose, Spruce, and Wisteria apply the recom
mended rates indicated below.

Insects	Amount EAGLE DIAZINON to use	
	per 3 gals. water	per 100 gals. water
Aphids		
Beet beetles		
Carnation bud moths		
Carnation shoot moths		
Crow mites		
Cyathidius mites		
Distortus leafminers		
European vine shoot moths		
European red mites		
Flea beetles		
Holly bud moths		
Leafhoppers	1 fl. oz. *	3 pt.
Obscure root weevils		
Omnivorous leafhoppers		
Prickly mites		
Scale crawlers		
Cottony cushion		
Lecanium		
Pine needle		
San Jose scale		
Thrips		
Two spotted mites		
Webworms		
Whiteflies		
Apple and thorn skeletonizers		
Cottonweevil webworms		
Fall webworms		
Hemlock chionias	1 1/2 fl. oz. *	3 pts. *
Oak looper		
Oak leaved sawfly		
Pink hoppers		
Tree caterpillars		

* 1 fl. oz. = 2 tbsp.

16 fl. oz. = 1 gal.

Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pine, and Gorse. A nice injury to the plants may occur. *Mimosa pudica* — Mix 1 qt oil (1 tablespoon) in 3 gal. of water (or 1 qt in 100 gal. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

To control insects such as Ants, Earwigs, Flies, etc., which are common pests, apply the product as a spray or dust to the areas where they enter the house, especially the kitchen and the bathroom. Repeat application if necessary.

Boxelder Bugs - 50
oz. (2 1/2-3 tablespoons)
toss thoroughly at
trunks, and into crevices.

LOWE'S CHURCH Bldg

Art. Armyed vs.
Omer Miller, Ex. 104
Columbus, Cuckers
Cuckers, Diner
Cuckers, Ex. 104

First Loan to: Duke S.
Washington, D.C.
Embassy

Baroness Olga A. von
Muller, Chicago
Finn. Lutheran

Phosphogate School

Thoroughly water
tion.
Satur extracts the
description and is r
ished when used in

...of the ...
...of ...
...of ...
...of ...
...of ...

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE