

38621-16



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)
CAUTION**

Keep out of reach of children. Harmful to some birds. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.
IF INHALED — Remove victim to fresh air. Apply respiration as indicated.
IF ON SKIN — Remove contaminated clothing and wash affected areas with soap and water.
IF ON EYES — Flush eyes for at least 15 minutes with water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS
This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop. Avoid crop or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
Pesticide Disposal — Pesticides that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
Containers Disposal — Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.
Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk harrow, or other suitable means for greatest effectiveness.

DIRECTIONS FOR USE CONTINUED ON BACK PANEL

**EAGLE
DYZON 14G
GRANULAR
INSECTICIDE**

For control of certain soil insects including wireworms, cutworms and other crop pests listed.

ACTIVE INGREDIENT:
O, O-diethyl O-(2-sopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%
INERT INGREDIENTS: 85.7%
TOTAL 100.0%

BEST DOCUMENT AVAILABLE

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

ACCEPTED

MARK 13 1974

**MANUFACTURED BY:
FOSTER MANUFACTURING, INC.
157 FIRST STREET, COACHELLA, CALIFORNIA 92230
(714) 398-6154**

EPA Reg. No. 38621-16 EPA Est. No. 11017-CA-01
NET WEIGHT: 50 POUNDS

**DIRECTIONS FOR USE, CONTINUED
FIELD AND FORAGE USE**

Soybeans (Plots)
Lesser Greenleaf Borers: Apply 7-14 lbs per acre in a 10-inch band centered over the row when beans are in the 2nd or 3rd trifoliate leaf stage.
Corn (Production Areas)
Lesser Greenleaf Borers: Apply 7-14 lbs per acre in a 10-inch band centered over the row at first signs of rootworms, but not later than the 4th leaf stage of the corn plant. Repeat application if necessary.
Lesser Greenleaf Borers: Apply 7-14 lbs per acre in a 10-inch band centered over the row at planting time or when crop begins to emerge. Repeat application if necessary.
Apply amount per acre recommended in the chart for basal treatment.

Suggested Rates According to Corn Row Spacing

Row Spacing	Linear feet of corn row per acre
45"	10,690
42"	12,476
40"	13,008
38"	13,959
36"	14,520
35"	17,474
28"	21,700
20"	26,136
15"	34,848

Note: For row spacings not listed, calculate dosage rates based on the 40-inch row spacing.

European Corn Borers: Apply 7-14 lbs per acre in a 10-inch band centered over the row when 75% of the corn plants are in the 4th leaf stage. Repeat application if necessary.
Southern Corn Rootworms: Apply 7-14 lbs per acre in a 10-inch band centered over the row when rootworms first appear. Repeat application if necessary.
Note: Do not feed treated forage to sheep for 10 days following application. Immediately following application, temporarily soaking of forage is recommended.
Cutworms: Broadcast 14-28 lbs per acre just prior to planting or emergence and work into soil 1-2 inches. Do not broadcast cutworms; apply just prior to planting and work into soil 1-2 inches.



BEST DOCUMENT AVAILABLE

38621-16

14G LAR ICIDE

insects including
er crop pests listed
DIENT:
(1)yl 4
14.3%
85.7%
TOTAL 100.0%

LABE
OF CHILDREN
ON

ADDITIONAL
STATEMENTS
TED

ED BY
YORING, INC.
LA, CALIFORNIA 92238
38621-16
Est. No. 11017-CA-01
0 POUNDS

DIRECTIONS FOR USE, CONTINUED FIELD AND FORAGE INSECTS

Beans (Pinto)
Lesser Cornstalk Borers Apply 7.14 lbs per acre...
Corn
Corn Rootworm Larvae...
Lesser Cornstalk Borers Apply 7.14 lbs per acre...
Apply amount per acre...
Suggested Rates According to Corn Row Spacing

Row Spacing	Amount of EAGLE DYZON 14G per Acre	Amount of EAGLE DYZON 14G per Acre
48	10.80	3.60
42	12.45	4.15
36	15.12	5.04
30	17.80	5.93
24	21.24	7.08
20	25.20	8.40
15	34.56	11.36

Note: For the spacing... dosage rates based on the 48 inch spacing.


European Corn Borers Apply 7.14 lbs per acre...
Fat Armyworms Apply 7.14 lbs per acre...
Southwestern Corn Borers Apply 7.14 lbs per acre...
Cutworms Broadcast 14.28 lbs per acre...

Seed Corn Maggots
Garden Symphylans (Centipedes)
Wireworms
Peanuts
Southern Corn Rootworms
Cutworms A
Lesser Cornstalk Borers
Sorghum
Cutworms B
Lesser Cornstalk Borers
Nymphs of the European Corn Borer
Soybeans
Cutworms C
Lesser Cornstalk Borers
Sugar Beets
Sugar Beet Root Maggots
Sugarcane
Sugarcane Borers
Wireworms
Cutworms D

Tobacco
Cutworms
Mole Crickets
Seed Corn Maggots
Wireworms
Forage Crops (Alfalfa, Clover, Coarse)
Cutworms
VEGETABLE CROP IN:
Broccoli, Brussels Sprouts, Cabbage
Root Maggots
Onions
Onion Maggots
Sweet Corn
Tomatoes
Vinegar Flies
Radishes
Root Maggots
OTHER VEGETABLE IN:
Cutworms (Surface and Subterranean)
Brussels Sprouts
Cabbage
Cauliflower
Cucumbers
Eggplant
Lettuce
Peas
Peppers
Potatoes
Garden Symphylans (Centipedes)
Cabbage
Carrots
Lettuce
Peas
Snap Beans
Mole Crickets
Broccoli
Brussels Sprouts
Cabbage
Cauliflower
Cucumbers

BEST DOCUMENT AVAILABLE

38621-16



Broadcast 10 to 12 lbs. per acre
 Broadcast 12 to 14 lbs. per acre
 Broadcast 14 to 16 lbs. per acre
 Broadcast 16 to 18 lbs. per acre
 Broadcast 18 to 20 lbs. per acre
 Broadcast 20 to 22 lbs. per acre
 Broadcast 22 to 24 lbs. per acre
 Broadcast 24 to 26 lbs. per acre
 Broadcast 26 to 28 lbs. per acre
 Broadcast 28 to 30 lbs. per acre
 Broadcast 30 to 32 lbs. per acre
 Broadcast 32 to 34 lbs. per acre
 Broadcast 34 to 36 lbs. per acre
 Broadcast 36 to 38 lbs. per acre
 Broadcast 38 to 40 lbs. per acre
 Broadcast 40 to 42 lbs. per acre
 Broadcast 42 to 44 lbs. per acre
 Broadcast 44 to 46 lbs. per acre
 Broadcast 46 to 48 lbs. per acre
 Broadcast 48 to 50 lbs. per acre

Chago
Cutworms
Mole Crickets
Wireworms
Field Corn Maggots
Winged Weevils
Forage Crops (Alfalfa, Clover, Concess and Lespedeza)
Cutworms
Wireworms

VEGETABLE CROP INSECTS
Broccoli, Brussels Sprouts, Cabbage and Cauliflower
Root Maggots
Wireworms
Onions
Onion Maggots
Sweet Corn
Tomatoes
Vinegar Flies (Drosophila spp.)
Radishes
Root Maggots

OTHER VEGETABLE INSECTS
Cutworms (Surface and Subterranean)
Beetle
Brussels Sprouts
Cabbage
Carrot
Cauliflower
Collards
Courtyards
Broccoli
Onions
Potatoes
Garden Symphylans (centipedes)
Cabbage
Carrots
Lettuce
Pars
Broccoli
Onions
Potatoes
Garden Symphylans (centipedes)
Cabbage
Carrots
Lettuce
Pars
Broccoli
Onions
Potatoes

Leafhopper
Wireworms
Onion
Onion Maggots
Sweet Corn
Tomatoes
Vinegar Flies (Drosophila spp.)
Radishes
Root Maggots

SMALL FRUIT INSECTS
Strawberries
Mole Crickets

LAWN INSECTS
Lawn Chinch Bugs
Beetles
White Grubs
Hydrocodon Beetles
Grasshoppers

MANUFACTURED BY
FOSTER MANUFACTURING, INC.
 1577 FIRST STREET, COACHELLA, CALIFORNIA 92226
 (71) 8-0124



BEST DOCUMENT AVAILABLE