



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(& DOMESTIC ANIMALS)**

**CAUTION**

Wash hands after use. May cause eye irritation. Avoid contact with skin or eyes. In case of eye contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists.

**ENVIRONMENTAL HAZARDS**

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinsate that cannot be used or chemically re-processed should be disposed of according to procedures approved by Federal, State or Local disposal authorities.

**CONTAINER DISPOSAL:** Consult Federal, State or Local disposal authorities for approved procedures.

Applications of materials containing sulphur should not be made prior to or during periods of excessively high summer temperatures or within 2 weeks of an oil spray. On citrus do not make an oil application within 60 days prior to or after a sulphur application.

**MIXING DIRECTIONS:** Fill the spray tank with water until the agitator blades are covered. With the agitator running sift in the required amount of EAGLE WETTABLE SULPHUR. Continue filling and keep agitator running.

**ALMONDS:** Brown Rot, Scab — Use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.



**EAGLE  
WETTABLE  
SULPHUR  
FUNGICIDE-  
INSECTICIDE**

<b>ACTIVE INGREDIENT:</b>	
Sulphur .....	92.0%
<b>INERT INGREDIENTS:</b> .....	9.0%
<b>TOTAL</b> .....	100.0%

This material in this bag is guaranteed to pass 95% through a 325 mesh screen.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS**

EPA Reg. No. 38621-17-AA

EPA Est. #11017-CA-01

MANUFACTURED BY:

**FOSTER MANUFACTURING CO.  
1577 FIRST STREET  
COACHELLA, CALIFORNIA 92236  
(714) 398-6154**

**NET CONTENTS:  
50 POUNDS**

**APPLES AND PEARS:** For Scab Mildew control, use as a spray in Calyx and Early cover sprays. Use 5 pounds of EAGLE WETTABLE SULPHUR or 5 pounds Polysulphide Compound or 5 gallons Liquid Lime Sulphur Solution per 100 gallons dilute spray. Sulphur may injure certain varieties of APPLES. Check specific variety sulphur prior to using. Pears only. Use after fruit begins to turn down. 100 lbs. per acre. Spray: Blister Mites, European Red Mite, Twospotted Spider Mites — Use 5 pounds of EAGLE WETTABLE SULPHUR plus 5 pounds Polysulphide Compound or 5 gallons Lime Sulphur Solution to each 100 gallons dilute spray.

**BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES:** Powdery Mildew — Use 5 pounds per 100 gallons of water. Apply when disease first appears and repeat at 7 to 10 day intervals.

**CHERRIES:** Brown Rot, Leaf Spot, Powdery Mildew, Rust, Scab and Stem Blight — Use 5 pounds per 100 gallons of water. (Apply a minimum of 200 gallons of water per acre.) Apply in the pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**CITRUS:** Citrus Flat Mite, Yuma Spotted Mite and Citrus Rust Mite — Use 5 pounds per 100 gallons of water. Apply when mites are present and repeat as necessary.

**GRAPES:** Powdery Mildew — Use 5 pounds per 100 gallons of dilute spray. Apply beginning when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals. Application preferred. Concord and type grapes may be injured by sulphur.

**PEACHES AND NECTARINES:** European Red Mite, Powdery Mildew, Rust, Scab — Use 5 to 10 pounds per 100 gallons (By ground apply a minimum of 500 gallons of water per acre.) Apply in the pre-bloom, bloom and post-bloom periods.



