

RECEIVED
APR 13 1981
Under the Federal
Fungicide, and Insecticide Act,
as amended, for the
registered under
EPA Reg. No. 38621-18



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS**

CAUTION

Avoid contact with skin, eyes, or clothing. Avoid breathing vapors, dust or spray mist. May be harmful if swallowed.

ENVIRONMENTAL HAZARDS

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open burning and dumping prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinseate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local disposal authorities.

CONTAINER DISPOSAL: Consult Federal, State or Local disposal authorities for approved procedures.

Make thorough, full coverage applications beginning when insects first appear and repeat as necessary to maintain control. May be applied by ground or air.

Applications of dusting and spraying materials containing sulphur should not be made prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On citrus do not make an oil application within 60 days prior to or after a sulphur application.

APPLES AND PEARS: Scab, **ALMONDS, CHERRIES, PEACHES, NECTARINES, PLUMS, PRUNES:** Brown Rot, Cherry Leaf Spot (Cherries only), Peach Silver Mite and Rust, **APPLIES, NECTARINES, PEACHES:** Powdery Mildew —

**EAGLE
DUSTING
SULPHUR
FUNGICIDE-
INSECTICIDE**



ACTIVE INGREDIENT:	
Sulphur	98.0%
INERT INGREDIENTS:	2.0%
TOTAL	100.0%

This material in this bag is guaranteed to pass 95% through a 325 mesh screen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 38621-18 EPA Est. #11017-CA-01

MANUFACTURED BY:
FOSTER MANUFACTURING INC.
1577 FIRST STREET
COACHELLA, CALIFORNIA 92236
(714) 398-6154

**NET CONTENTS:
50 POUNDS**

4.3



38621-18



Use 40 to 50 pounds per acre in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. On Pears do not apply after the fruit begins to turn down. Sulphur may injure certain varieties of apples, check tolerance prior to using.

GRAPES: Powdery Mildew — Use 8 to 15 Pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat before blossoms open and continue at 10-14 day intervals as necessary.

BLACKBERRIES, BLUEBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES, CURRANTS: Powdery Mildew — Use 10 1/4 to 20 pounds per acre. Apply when disease first appears and repeat at 7-10 day intervals.

FIG: Fig Rust Mite. 40 pounds per acre in the spring and again in the early summer.

PLUM, PRUNE: Brown Rot, Prune Rot, 30 to 35 pounds per acre 5 weeks before harvest. Repeat each 7 days if wet weather prevails during the period prior to harvest.

CITRUS: Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite and Citrus Rust Mite — Use 100 pounds per acre. Apply when pests first appear and repeat as necessary.

VEGETABLE AND FIELD CROPS: Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

BEANS: Powdery Mildew, Rust, Thrips, Spider Mites — Apply 15 to 30 pounds per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

CARROTS: Petrobia Mite — Apply 30 to 35 pounds per acre.

COTTON: Strawberry Spider Mite — Use 35 to 40 pounds per acre when mites first appear and repeat as necessary.

DATES: To control Banks Grass Mites use 25 to 50 pounds per acre, applying by ground or air. Use the higher rates by air. Thorough coverage is essential for good control.

MELONS: Powdery Mildew. 15 to 20 pounds per acre. Spider Mite 20 to 40 pounds per acre. Do not apply to varieties susceptible to sulfur injury.

ONIONS, GARLIC: Petrobia Mite — Apply 26 to 30 pounds per acre.

PEAS, TURNIPS, RUTABAGAS: Powdery Mildew. 15 to 20 pounds per acre.

SQUASH: (Summer or Winter): Powdery Mildew, 15 to 20 pounds per acre. Begin when first true leaves appear. Repeat at weekly intervals. Do not apply to varieties susceptible to sulfur injury.

SUGAR BEET: Spider Mite, 46 to 50 pounds per acre.

ALFALFA (grown for seed): Atlantic Mite, 15 to 40 pounds per acre when first observed in flower heads and before defoliation of more than the bottom leaves has occurred. Repeat as needed throughout seed-set. May cause some flower burn.

TOMATOES: Tomato Russet Mite — Apply 30 to 40 pounds per acre.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



43