

PRECAUTIONARY STATEMENTS
Hazards to Humans and
Domestic Animals

CAUTION

Harmful if swallowed. Avoid contact with eyes, contact with skin, inhalation of mist and contamination of food and feed.

ENVIRONMENTAL HAZARDS

Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a full type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered until sprays have dried. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information.

WARNING

Area treated with Prometryne (insert date and time). Do not enter until spray dries (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

Storage: Do not contaminate water, food, or feed by storage or disposal. Store in a dry place.

Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

In case of minor spills, follow all precautions indicated above and clean-up immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

GENERAL INFORMATION

Prometrex 4L is a selective herbicide that may be applied either before or after weeds emerge for control of most annual broadleaf weeds and grasses, including pigweed (carelessweed), Florida pusley, smartweed, teaweed (prickly sida), annual morningglory, ragweed, groundcherry, mustard, lambsquarters, purslane, malva, junglerice, foxtail, wild oats, barnyardgrass (watergrass), crabgrass, signalgrass, and other Brachiaria spp., Panicum spp., and goosegrass. Prometryne also controls shallow germinating seedlings of cocklebur, colleweed, and sandbur. It does not control johnsongrass, bermudagrass, other established perennials, or sprangletop at selective rates.

When applied before weeds emerge, Prometryne enters weeds through their roots. Thus, its effectiveness depends on rainfall or irrigation to move it into the soil. Very dry soil conditions following applica-

PROMETREX

Herbicide for selective weed control in

ACTIVE INGREDIENT:

Prometryne: 2,4-bis (isopropyl-amino)-6-(methyl-thio)-s-triazine

INERT INGREDIENTS:

CONTAINS 4#/GAL. A.L.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Statement of Practical Treatment

IF ON SKIN—Wash with plenty of soap and water until irritation persists.

IF IN EYES—Flush with water. Get medical attention immediately.

See side panel for additional precautionary statements.

WARRANTY STATEMENT

NOTICE: Seller warrants that the product conforms to its label directions under normal conditions of use, but neither this warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE nor implied, extends to the use of this product under abnormal conditions or under conditions not reasonably foreseeable. Buyer assumes the risk of any such use.

Manufactured for:

MAKHTESHIM-AGAN (AMERICA) INC

2 Park Avenue
New York, New York

ACCEPTED

3 1 3 1986

Insecticide,
Fungicide, or
Herbicide Act,
Label for pesticide

38652-4

EPA Reg. No. 38652-

EPA Est. No. 11603-I

Mixing: (1) Fill spray tank 1/2 to 3/4 full with clean water. (2) Start agitation. (3) Pour the product directly from the container into the spray tank partially filled with water, and then add the rest of the water. (4) Provide agitation during mixing and application to maintain a uniform suspension.

CLEANING EQUIPMENT

Wash sprayer thoroughly with clean water immediately after use. Sprayers used to apply Prometrex 4L should not be used to apply pesticides on sensitive crops such as tobacco, sugar beets, or vegetables, as even small residues of Prometryne in the tank may cause injury to these crops.

SEEDBED PREPARATION

To insure proper placement of Prometrex 4L seedbeds must be

EX 4L

control in cotton and celery

44.0%

56.0%

TOTAL 100%

UTION

CH OF CHILDREN

ctical Treatment
ap and water. Get medical attention if
medical attention if irritation persists.
Precautionary statements.

STATEMENT

...forms to its chemical description and is
the label when used in accordance with
but neither this warranty nor any other war-
FOR A PARTICULAR PURPOSE, express
just contrary to label instruction or under
not reasonably foreseeable to seller, and

AMERICA) INC.

ew York 10016

38652-
11603-IS-01

PREPLANT INCORPORATION (Arizona, California, and New Mexico).
Apply Prometrex 4L as a broadcast or band treatment. If broad-
cast, treat the flat soil surface prior to listing. If banded, apply over par-
tially finished or finished beds. Incorporate up to 4 inches deep im-
mediately after application with PTO-driven equipment, double disk,
rolling cultivator, or bed conditioner.

PREPLANT

RATES OF APPLICATION

REGION	SOIL TEXTURE	BROADCAST* RATE PER ACRE
	sand, loamy sand	Do not use
Arizona, California, and New Mexico	sandy loam, loam	4 pts.
	silt loam, clay	6 pts.

*For band treatment, the amount of Prometrex 4L required may be
calculated by the formula

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

Precautions: Do not use Prometrex 4L on sand or loamy
sand soils, in cut areas of newly leveled fields, in areas of excess
silt (excess of ECE values of 1.0), or in areas where flooding
over the bed is likely to occur as crop injury may result. Do not
plant cotton in tractor wheel depressions as crop injury may
result. On mulch planted cotton, water back only after cotton
seedlings are well established. In New Mexico, apply either pre-
plant incorporated or preemergence (not both)—see Pre-
emergence section.

PREEMERGENCE

Apply at planting or shortly after planting. Prometrex 4L may be
used on cotton planted flat, on beds, or in furrows. To avoid concentra-
tion of Prometrex 4L in the seed furrow, do not make broadcast
applications to cotton planted in furrows more than two inches deep.
Band applications may be made to cotton planted in furrows deeper
than two inches, but band width should not exceed the width of the bot-
tom of the furrow. Do not use on shallow soils with caliche subsoils or in
areas with caliche out-croppings.

Rainfall or irrigation is needed following preemergence application to
obtain weed control.

If necessary, cotton may be replanted in soil previously treated with
Prometrex 4L. Do not apply a second preemergence application of
Prometryne as injury may occur.

PREEMERGENCE (Do not use on sand or loamy sand soil).

Rates of Application

Region	Soil texture	Broadcast* rate per acre
Mid-South & Southeast other than Mississippi River Delta in Miss.	sandy loam	4-5 pts.
	silt and clay loam	6 pts.
	Sharkey Clay (Arkansas only)	7 pts.
Mississippi River Delta in Miss.	sandy loam	5-6 pts.
	silt and clay loam	7 pts.
	Sharkey clay	Do not use
Blacklands of Texas & Okla. Texas Gulf Coast, & Texas Coastal Bend	loam	3 pts.
	clay	6 pts.
Rio Grande Valley of Texas**	loam	4 pts.
	clay	6 pts.
High Plains, Rolling Plains and Edwards Plateau of Texas, Southwest Texas and New Mexico***	sand loamy sand	Do not use
	sandy loam	2 pts.
	loam, sandy clay loam	3 pts.
	other clay soils	4 pts.
Arizona and California		Do not use

*For band treatment, the amount of Prometrex 4L required may be
calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

Do not cover treated bands with soil while cultivating untreated row

or use.
to apply
or vege-
ause in-

must be

New Mexico)
ont. If broad
ply over par
res deep im-
double disk.

POSTEMERGENCE DIRECTED

Take special care in applying Prometrex 4L postemergence to prevent contact of the spray with cotton leaves, or injury may occur. Recommended practices include (1) using precision application equipment so the spray is accurately directed to the base of the cotton plants and still thoroughly covers the soil and weeds beneath the cotton plants, (2) applying during calm periods to prevent drift, (3) using branch lifters or shields if leaf contact cannot be avoided merely by directing the spray, (4) applying only when all plants have exceeded the minimum recommended height, and (5) applying to level, well-prepared surfaces such as beds made with bed-shapers that are relatively free of clods.

Apply to cotton less than 10 inches tall only if cotton is bed or flat-planted. Do not apply to furrow-planted cotton until furrows are leveled (plowed in).

When applying to emerged weeds, add 2 qts. of surfactant per 100 gals. of spray mixture. Use a surfactant that is compatible with Prometrex 4L when applied in cotton and is approved by EPA for use on food and feed crops. Examples include X-77, Tronic, and Trilon.

Do not treat cotton under stress from drought, cultivator damage, or fertilizer application.

Cotton three to six inches tall (Texas and the Mid-South (Arkansas, Louisiana, Mississippi, Missouri, and Tennessee) only): Take extreme care in applying Prometrex 4L to cotton three to six inches tall so the spray does not contact cotton leaves. Apply with a precision applicator equipped with fenders or shields, such as Bell Row Shield, Dickey Fenders, or W & A Fenders.

Cotton six or more inches tall (all regions): Applications of Prometrex 4L may be made any time after cotton is six inches tall.

To Control Emerged Weeds Only (Chemical Hoe): All soil textures, two or three applications may be necessary.

Height of cotton and area of use	Height of weeds	Broadcast* rate per acre
3 to 6 inches Texas and the Mid-South (Arkansas, Louisiana, Mississippi, Missouri and Tennessee)	less than 1 inch	1 1/4 pts.
6 or more inches All regions	less than 2 inches	1 1/2 - 1 3/4 pts.

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

To Control Emerged Weeds and Germinating Weeds (Layby): One application per season

Region*	Soil Texture	Broadcast rate/acre	Height of Cotton	Height of Weeds
Mid-South and Southeast	sandy	3 pts.	at least 2 inches	less than 2 inches
	loam	3 1/2 pts.		
	silt loam	4 pts.	at least 12 inches	less than 2 inches
Blacklands of Texas and Oklahoma	clay	4 pts.		
High Plains of Texas and New Mexico	sandy	2 pts.	at least 12 inches	less than 2 inches
	loam and clay	3 pts.		
Southwest Texas	loam	3 pts.	at least 12 inches	less than 2 inches
	clay	4 pts.		
Rio Grande Valley of Texas		Do not use		
Arizona and California (Do not use in the Coachella Valley)	sand and loamy sand	Do not use		
	loam	4 pts.	at least 12 inches (or 18 inches where flood type nozzles are used)	less than 2 inches

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

ROTATIONAL CROPS
The vegetable crops and cover crops listed below may be planted in

CELERY
POSTEMERGENCE TREATMENT OF SEEDBEDS (Florida Broadcast 1 1/2 - 2 pts. in 20-40 gals. of water per acre after celery true leaves. Application may be made over the growing celery only after seedbed covers have been removed from seedbeds at least one week. Apply only once per year to seedbeds.

DIRECT-SEEDED CELERY (California only)
Apply Prometrex 4L at rates given below in a minimum of 20 gals. of water per acre. Within the rate ranges given, use the lower rate on coarse-textured soils and soils low in organic matter, use the higher rate on fine-textured soils and soils high in organic matter.

Preemergence: Broadcast 3-4 pts. per acre at planting or shortly before planting before celery emerges.

Postemergence: Broadcast 2-2 1/2 pts. per acre after celery true leaves. Application may be made over the growing celery before weeds are 2 inches tall.

To Avoid Injury to Direct-seeded Celery: (1) Make either preemergence or one postemergence application (not both) to celery crop. (2) Do not use on sand or loamy sand. (3) Do not apply if celery is under water stress. (4) Do not apply postemergence treatments of Prometrex 4L with other pesticides. Apply only after foliar applications or other pesticides are dry. (5) Do not apply within two weeks after an application of herbicidal oil, such as "carrot" oil.

POSTEMERGENCE TREATMENT OF TRANSPLANTS
Apply one application at the appropriate rate from the following a minimum of 20 gals. of water per acre during the 2-6 weeks after transplanting. Within the rate ranges given, use the lower rate on relatively coarse-textured soils and soils low in organic matter, use the higher rate on relatively fine-textured soils and soils high in organic matter. Application may be made over the growing celery. Apply before weeds are 2 inches tall.

State	Soil	Broadcast rate per acre
Florida	sandy or muck	2-4
California	coarse-textured	2 1/2 - 4
	fine-textured	4-5
Hawaii	coarse-textured	4-6
	fine-textured	6-8

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

ROTATIONAL CROPS
The following crops may be seeded 5 months after applying more than 3 pts. Prometrex 4L per acre on celery: cabbage, corn, okra, onions, peas, and red beets.

1ST*
ACRE

ired may be

or loamy
excess
flooding
Do not
ary may
r cotton
her pre-
ee Pre-

4L may be
concentra-
broadcast
ches deep.
as deeper
of the bot-
soils or in

ication to

reated with
plication of

oil).

least*
per acre

pts.

ts.

ts.

pts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

ts.

expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered until sprays have dried. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information.

WARNING

Area treated with Prometryne (Insert date and time). Do not enter until spray dries (Insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

Storage: Do not contaminate water, food, or feed by storage or disposal. Store in a dry place.

Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

In case of minor spills, follow all precautions indicated above and clean-up immediately. Sweep up and dispose of wastes and broken or empty containers in a landfill approved for pesticide use.

GENERAL INFORMATION

Prometrex 4L is a selective herbicide that may be applied either before or after weeds emerge for control of most annual broadleaf weeds and grasses, including pigweed (carelessweed), Florida pusley, smartweed, leaved prickly sida, annual morningglory, ragweed, groundcherry, mustard, lambsquarters, purslane, malva, junglerice, foxtail, wild oats, barnyardgrass (watergrass), crabgrass, signalgrass, and other Brachiaria spp., Panicum spp., and goosegrass. Prometryne also controls shallow germinating seedlings of cocklebur, coffee-weed, and sandbur. It does not control johnsongrass, bermudagrass, other established perennials, or sprangletop at selective rates.

When applied before weeds emerge, Prometryne enters weeds through their roots. Thus, its effectiveness depends on rainfall or irrigation to move it into the soil. Very dry soil conditions following application may result in poor weed control. Under these conditions a shallow cultivation or rotary hoeing will generally result in better weed control.

When applied to emerged weeds as a directed spray, Prometryne provides foliar knockdown of existing weeds and/or residual control of later germinating weeds, depending on the rate applied.

Equipment

Use conventional spray equipment with hydraulic or mechanical agitation. Screens in nozzles and in suction and in-line strainers should be no finer than 50-mesh. Use a pump with capacity to maintain 35-40 psi at the nozzles. If hydraulic agitation is used, the pump should also provide sufficient agitation in the tank to keep the mixture in suspension.

Nozzles: For preplant incorporated or preemergence application use fan-type nozzle tips. For postemergence band application use off-center nozzle tips. For postemergence broadcast application use fan-type nozzle tips or off-center nozzle tips. Use flood-type nozzle tips only in Arizona and California for making layby treatments to cotton at least 18 inches tall.

Statement of Practical Treatment

IF ON SKIN—Wash with plenty of soap and water until irritation persists.

IF IN EYES—Flush with water. Get medical attention.

See side panel for additional precautions.

WARRANTY STATEMENT

NOTICE: Seller warrants that the product conforms to its reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or implied, extends to the use of this product under abnormal conditions or under conditions not reasonably foreseeable. Buyer assumes the risk of any such use.

Manufactured for:

MAKHTESHIM-AGAN (AMERICA) INC.

2 Park Avenue
New York, New York

ACCEPTED

3 1 3 1986

Insecticide,
Fungicide,
and Herbicide Act,
EPA Reg. No. 38652-4

EPA Reg. No. 38652-

EPA Est No. 11603-

Mixing: (1) Fill spray tank 1/2 to 3/4 full with clean water. (2) Start agitation. (3) Pour the product directly from the container into the spray tank partially filled with water, and then add the rest of the water. (4) Provide agitation during mixing and application to maintain a uniform suspension.

CLEANING EQUIPMENT

Wash sprayer thoroughly with clean water immediately after use. Sprayers used to apply Prometrex 4L should not be used to apply pesticides on sensitive crops such as tobacco, sugar beets, or vegetables, as even small residues of Prometryne in the tank may cause injury to these crops.

SEEDBED PREPARATION

To insure proper placement of Prometrex 4L seedbeds must be well prepared and as free as possible from trash and clods. A firm seedbed is best for obtaining effective weed control. Uniformity in height and width of seedbed is essential for proper postemergence applications of Prometrex 4L. Beds should be low and flat. Take care to avoid planter marks. Wide plant packer wheels or rollers are recommended. Wheel furrows should also be uniform in depth. Mount the sprayer so that it follows the same rows as the planter.

COTTON

PROMETREX 4L may be applied preplant incorporated or preemergence and/or postemergence as recommended in the following tables. The postemergence applications may follow preplant incorporated or preemergence treatments of Prometrex 4L.

Apply Prometrex 4L in 20-40 gals. of water per acre as a broadcast or band application at the rates indicated in the following tables.

Do not use on glandless cotton varieties, as crop injury will occur.

NOTE: Do not feed treated forage to livestock, or graze treated areas.

ctical Treatment
 ap and water. Get medical attention if
 medical attention if irritation persists.
 / precautionary statements.

STATEMENT
 conforms to its chemical description and is
 the label when used in accordance with
 but neither this warranty nor any other war-
 FOR A PARTICULAR PURPOSE, express
 duct, contrary to label instruction or under
 not reasonably foreseeable to seller, and

AMERICA) INC.
 ew York 10016

38652-
 11603-IS-01

er use.
 to apply
 or vege-
 ause in-

must be
 . A firm
 mply in
 nce ap-
 ke care
 recom-
 unt the

led or
 follow-
 incor-

oadcast
 es.
 cur.
 areas.

seedlings are well established. In New Mexico, apply either pre-
 plant incorporated or preemergence (not both)—see Pre-
 emergence section.

PREEMERGENCE
 Apply at planting or shortly after planting Prometryn 4L may be
 used on cotton planted flat, on beds, or in furrows. To avoid concentra-
 tion of Prometryn 4L in the seed furrow, do not make broadcast
 applications to cotton planted in furrows more than two inches deep.
 Band applications may be made to cotton planted in furrows deeper
 than two inches, but band width should not exceed the width of the bot-
 tom of the furrow. Do not use on shallow soils with caliche subsoils or in
 areas with caliche outcroppings.

Rainfall or irrigation is needed following preemergence application to
 obtain weed control.

If necessary, cotton may be replanted in soil previously treated with
 Prometryn 4L. Do not apply a second preemergence application of
 Prometryn as injury may occur.

PREEMERGENCE (Do not use on sand or loamy sand soil).

Rates of Application

Region	Soil texture	Broadcast* rate per acre
Mid-South & Southeast other than Mississippi River Delta in Miss.	sandy loam	4-5 pts.
	silt and clay loam	6 pts.
	Sharkey Clay (Arkansas only)	7 pts.
Mississippi River Delta in Miss.	sandy loam	5-6 pts.
	silt and clay loam	7 pts.
	Sharkey clay	Do not use
Blacklands of Texas & Okla. Texas Gulf Coast, & Texas Coastal Bend	loam	3 pts.
	clay	6 pts.
Rio Grande Valley of Texas**	loam	4 pts.
	clay	6 pts.
High Plains, Rolling Plains and Edwards Plateau of Texas, Southwest Texas and New Mexico***	sand loamy sand	Do not use
	sandy loam	2 pts.
Arizona and California	loam, sandy clay loam	3 pts.
	other clay soils	4 pts.
Arizona and California		Do not use

*For band treatment, the amount of Prometryn 4L required may be
 calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

Do not cover treated bands with soil while cultivating untreated row
 middles.

**Rio Grande Valley of Texas—furrow irrigated cotton—to insure good
 weed control, make a shallow cultivation if adequate rain does not fall
 soon after applicatic

***New Mexico—apply either preplant incorporated or preemergence
 (not both)—see Pre-plant Incorporation section.

of cotton
the pre-
ee Pre-

Height of cotton and area of use	Height of weeds	Broadcast rate per acre
3 to 6 inches Texas and the Mid-South (Arkansas, Louisiana, Mississippi Missouri and Tennessee)	less than 1 inch	1 1/4 pts.
6 or more inches All regions	less than 2 inches	1 1/4 - 1 3/4 pts

relatively coarse-textured soils and soils low in organic matter; higher rate on relatively fine-textured soils and soils high in matter. Application may be made over the growing celery. Applied weeds are 2 inches tall.

4L may be
concentra-
broadcast
ches deep.
as deeper
of the bot-
soils or in

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

ication to
reated with
ication of

To Control Emerged Weeds and Germinating Weeds (Layby): One application per season

Region	Soil Texture	Broadcast rate/acre	Height of Cotton	Height of Weeds
Mid-South and Southeast	sandy	3 pts.	at least 2 inches	less than 2 inches
	loam	3 1/2 pts.		
	clay loam	4 pts.		
Blacklands of Texas and Oklahoma	clay	4 pts.	at least 12 inches	less than 2 inches
High Plains of Texas and New Mexico	sandy	2 pts.	at least 12 inches	less than 2 inches
	loam and clay	3 pts.		
Southwest Texas	loam	3 pts.	at least 12 inches	less than 2 inches
	clay	4 pts.		
Rio Grande Valley of Texas		Do not use		
Arizona and California (Do not use in the Coachella Valley)	sand and loamy sand	Do not use		
	loam	4 pts.	at least 12 inches (or 18 inches where flood type nozzles are used)	less than 2 inches

State	Soil	Broadcast rate per acre
Florida	sandy or muck	2-4 p
California	coarse-textured	2 1/2 - 4
	fine-textured	4-5 p
Hawaii	coarse-textured	4-6 p
	fine-textured	6-8 p

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

oil).

Least
er acre

pts.

ts.

ts.

ts.

ts.

ts.

ts.

Use

ts.

ts.

Use

red may be

*For band treatment, the amount of Prometrex 4L required may be calculated by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed}$$

ROTATIONAL CROPS

The vegetable crops and cover crops listed below may be planted in the fall when Prometrex 4L was applied on cotton by no more than one of these methods that year preplant incorporated, preemergence, or only once at 1 1/4 - 1 3/4 pts. per acre postemergence.

Vegetables
cabbage, okra, onions, peas, red beets, sweet corn

Cover Crops (must be plowed down and not used for food or feed)
oats, sorghum, winter barley, winter rye, winter wheat.

Spring-seeded crops in California and Arizona and spring-seeded vegetables in the Rio Grande Valley of Texas should not be planted until after April 1.

reated row

sure good
es not fail

emergence