

38664-25 11 11

ACCEPTED
FEB 24 1986
Under the Federal In-
Fungicide, and Rodenticide
Act amended, for the purpose
of being used under
EPA Reg. No. 38664-2



PY-3 Fogger

Contains PYRENONE®
SUPER STRENGTH - Highly Active Solvent
System - Non-Aqueous
For Control of Residential, Industrial
and Garden Insect Pests.
Contains PYRENONE®
SUPER STRENGTH - Highly Active Solvent
System - Non-Aqueous
For Control of Residential, Industrial
and Garden Insect Pests.

ACTIVE INGREDIENTS:	
Pyrethrins.....	0.3%
*Piperonyl Butoxide, Technical.....	2.4%
Petroleum Distillate.....	1.5%
INERT INGREDIENTS:.....	95.8%
	100.00%

*Equivalent to 2% (butylcarbityl) (6-propylpipe-ronyl) ether and 0.4% related compounds.

PYRENONE - Registered trademark of Fairfield American Corporation.

EPA Reg. No. 38664-25 EPA Est. No. 38664-AR-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
Harmful if swallowed. Avoid breathing of vapor. Avoid contact with skin and eyes. See rear panel for additional precautions.

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION
Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid breathing of vapor. Avoid contact with skin and eyes.
When using this product as a space spray in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.
Do not remain in treated areas and ventilate the area after treatment is completed.
PHYSICAL OR CHEMICAL HAZARDS
Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Hold container upright with nozzle away from you. Press valve down and spray as directed. Keep nozzle at least 36 inches from surfaces or fabrics to avoid possible staining.
Remove pets, birds and cover fish aquariums before spraying.

FLIES, MOSQUITOES, GNATS, WASPS AND FLYING MOTHS: Close all doors and windows and spray room at the rate of 2-3 seconds per 1000 cu. ft. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Repeat treatment as necessary.

ROACHES, WATERBUGS, SILVERFISH, CRICKETS, SPIDERS, CENTIPEDES AND SOWBUGS: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

INDIAN MEAL MOTHS, ANGOUMOIS GRAIN MOTHS: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cu. ft. of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed.

Exposed stages of **CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, RED FLOUR BEETLES, RICE WEEVILS, GRANARY WEEVILS, GRAIN MITES:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 sq. ft. of surface. Keep area closed for 45 minutes to 1 hr. after treatment. Repeat applications as necessary.

CARPET BEETLES: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

BEDBUGS: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary.

FLEAS: Spray sleeping quarters, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

FOR THE CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese beetles hit by spray, as well as aphids, exposed thrips and spider mites.

May be used on plants, as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwood, English ivy, stonymus, tuschia, geraniums, crassula, grape vine, kentia palm, laurel, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against aphids, spider mites, exposed thrips, lace bugs, Japanese beetles, pavement ants, clover mites and army worms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Do not wet foliage or blossoms.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NET CONTENTS _____

DISPOSAL
Do not reuse empty container. Wrap container and put in trash collection.