



038733-00012-121583

Printed: 10:48:13 Friday, 28 May, 1999 # 76 / 3196

Systems Integration Group, Inc.



# DSMA LIQUID PLUS SURFACTANT

## DSMA Liquid For Selective Post-Emergent Weed Control

Contains 3.6 lbs. Disodium Methanearsonate Hexahydrate per Gallon

For Use In Cotton—Non-Crop & Ornamental Turf Areas  
For Grassy Weeds such as Johnson Grass and Certain Broadleaf Weeds  
such as Cocklebur and Other Weeds as Listed

### ACTIVE INGREDIENTS:

Disodium Acid Methanearsonate ..... 21.80%

INERT INGREDIENTS: ..... 78.20%

Total 100.00%

Total Arsenic (as elemental) all in water-soluble form ..... 8.90%

EPA Reg. No. 38733-12 EPA Est. No. 38733-FL-1

Keep Out of Reach of Children

## CAUTION

See Side Panel For Additional Precautionary Statements

### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed** — Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with fingers. Do not induce vomiting or give anything to an unconscious person.

**If In Eyes** — Flush with plenty of water and call a physician immediately.

**If On Skin** — Remove contaminated clothing. Wash immediately with soap and water. Call a physician immediately.

NET CONTENTS

GALLON

**BILCO ASSOCIATES**

AGRICULTURAL CHEMICALS  
ROBERT W. BILLETTE, INC. D-B-A BILCO ASSOCIATES

1140 West 50th Street • Hialeah, FL 33012

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**General Information:** This product is recommended for selective Post-Emergent weed control in Cotton, Non-Crop areas and for Lawn and Ornamental Turf. Its phytotoxic properties are quickly deactivated on contact with the soil. Spray when temperatures are warm (70°F +) and weeds are actively growing.

**Mixing:** This product must be thoroughly dissolved into the spray solution. Fill tank 1/2 full; add the recommended amount of herbicide as the agitator is operating and then continue to fill tank. Do not store solution in tank for a prolonged period (24 hours). Clean application equipment thoroughly after using by flushing with water in a safe place. (See Storage & Disposal Statement)

**Surfactant Spray Additive:** This product contains a surfactant.

**How to Apply:** Use a Low Pressure properly calibrated sprayer having adequate pumping and by-pass agitation. Adjust the Fan Type nozzles to give maximum coverage of target weeds with a minimum spray coverage of desirable plants or vegetation.

**Weeds Controlled - Grassy Weeds:** Johnson Grass, Nutsedge, Foxtail, Dallis Grass, Barnyard Grass, Sand Bur, Goose Grass, Bahia Grass, Smooth and Large Crabgrass, Broadleaf Weeds: Cocklebur, Puncture Vine, Chickweed, Wood Sorrel and Pigweed.

**Cotton:** Apply this product as a Post-Emergent spray to control weeds as listed. Keep spray off cotton foliage by "Directed" spray nozzles. Mix 1 Gallon of this product in 40 gallons of water to treat one acre as a "Directed" broadcast spray when weeds are small, 1-2 inches.

**For Band Application** (12" to 14" band on 40" planted rows): apply one gallon of the above diluted spray mixture for each 1 inch band width, per acre. Keep spray off cotton.

Band Width	Amt. Concentrate	Amt. Water/Acre
12"	38 1/2 Ounces	12 Gallons
14"	44 1/4 Ounces	14 Gallons

Application in Florida should be limited to Band Treatment only.

**When to Treat:** Treat when cotton is at least 3 inches high up until the first blooms appear. A second application, if necessary, should be applied 1 to 3 weeks after the first treatment. Do not apply after first bloom.

**Equipment:** Use off center fan type nozzles with 100 mesh screens, adjusted to 5 inches from the soil and spray at about 25 lbs. pressure.

**Note:** There may be a slight discoloration to Cotton plants at this recommended rate, however plants will develop normally and the yield will not be affected.

**Notice:** Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

Continued on other side panel

ACCEPTED

DEC 15 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 38733-12

Ornamental Turfgrass listed hereon. Observe use on new lawns until fully discolored without tolerant. Do not use or may be damaged.

Mow turfgrass to height water per 1,000 sq. ft. (all undesirable plants, intervals to give desired

This spray is particularly the lawn. These grass possible or at least le results will be obtained allow grass to dry before Use 3 ounces in a 2 g

**Non-Crop Areas:** This rights-of-way, fence row as listed hereon. Mix a

Spray plants until point is necessary for effective control.

PR  
Hazard

Harmful if swallowed. Contact with eyes. In case mist. Keep children and been washed into the

Do not feed treated for drift onto vegetables, weather conditions fa

STO

Do not contaminate w Pesticide, spray mixt ions must be dispose Resource Conservation lent) and offer for recy other approved State