

1/2 38832-1 PM-31

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460	EPA REGISTRATION NO. 38832-1	DATE OF ISSUANCE JUN 27 1989
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
 (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[] []

[] []

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

[Faint, mostly illegible text, likely a copy of the pesticide label or registration details.]

[Handwritten signature]
 []
 Product Manager (TS)
 Registration Division

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

WEST C-101

INDUSTRIAL MICROBICIDE

ACTIVE INGREDIENTS:

Methylene bis(thiocyanate)

2-Thiocyanomethylthiothiazole

INERT INGREDIENTS

This product weighs 7.99 pounds per gallon.

25%
25%
50%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive, causes irreversible eye damage and causes skin damage. Harmful or fatal if swallowed or absorbed through the skin. This product is a potential skin sensitizer. Workmen handling this product or treated material should wear impervious gloves, goggles or face shield, and protective clothing. All protective clothing, work shoes or boots, and equipment must be left at the work site at the end of the day. Eating, drinking, or smoking during use of this product is prohibited.

FIRST AID: In case of skin contact, wash promptly and thoroughly with soap and water, and finally with glycerin. If product gets in the eyes, flush immediately with copious amounts of clean, cool water for 15 to 30 minutes. Get medical attention immediately. If product is swallowed, call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water or wetlands. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: Do not expose to extreme temperatures.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not expose to extreme temperatures. Do not stack more than five drums high. Drums should be opened in well-ventilated areas. Leaking or damaged drums should be placed in overpack drums for disposal. Spills should be absorbed in sawdust or sand and disposed of in a sanitary landfill. Keep container closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at your EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with these directions.

COOLING TOWERS: WEST C-101

Wood rotting, soft or surface rot and internal or dry rot. It is effective against wood rotting fungi containing 2.0% to 2.8% WEST C-101

lakes. The amount applied should provide 2.4 to 3.2 lb of WEST C-101

1000 ft² of wood surface. Soft or surface rot can also be treated with WEST C-101 to the recirculating

water. The dosage should provide 5.0 lb of WEST C-101

water and the bleed-off should be stopped for 4 to 6 hr after treatment. Repeat every four months.

COOLING WATER: WEST C-101

and fungi in industrial recirculating cooling water systems should be cleaned thoroughly to remove old algal growth, microorganisms, and debris. The system should then be drained, flushed, refilled with water, and treated with WEST C-101 per 1000 gal of water.

of 0.8 to 4.8 fl oz per 1000 gal should be made every 1 to 5 days, depending on the severity of microbiological fouling.

DRILLING FLUIDS: To inhibit bacterial and fungal degradation of well fluids, WEST C-101 is used at concentrations of 0.2 to 1.0% based on the total wet weight of the fluid.

PETROLEUM SECONDARY RECOVERY: WEST C-101 is used to control sulfate-reducing bacteria, slime-forming bacteria, and other oil field water problems. WEST C-101 per 1000 gal of water.

made continuously or intermittently by means of a metering pump after injection pumps and injection well headers. Continuous flow treatment, add 15.6 to 52.0 fl oz WEST C-101 per 1000 gal of water.

Intermittent or Slug Method: When system is not treated, add 15.6 to 52.0 fl oz WEST C-101 per 1000 gal of water.

and 1 in 4 times per week, or as needed to maintain control.

CRUDE AND REFINED OILS: WEST C-101 is used as a biocide for the control of bacteria and fungi that cause the degradation of oil during storage. It should be added to the oil as it is being transferred to storage tank at the rate of 2.4 to 24.0 fl oz per 1000 gal of oil. Addition should be made batchwise where mixing occurs or transfer pump.

Unc
Fun
and
reg

Manufactured by:

WATER ENERGY SYSTEMS, INC.

980 EAST RAYMOND ROAD

FRUIT HEIGHTS, UTAH 84002

EPA REG. NO. _____ EPA EST. NO. _____

NET CONTENTS _____