KILLS: Ants, Brown Dog Ticks, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Scorpions, Silverfish, Sowbugs, and Spiders. Use outdoors against Fleas, Flies, Gnats, Mosquitoes, and Pantries.

Exposed Stages of Carpet, Cigarette, Drugstore, Indian Meal, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils.

RESTRICTIONS

"This Product is Toxic to Fish and Wildlife. Birds Feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label."

CAUTION

May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands and face thoroughly with soap and water before eating, drinking, or smoking. Wash all contaminated clothing with soap and hot water before reuse.

SHOULD ILLNESS OCCUR, GET IMMEDIATE MEDICAL AID

TO PHYSICIAN - Active Sulfide is Antidote

* Bayer is the registered trademark of Farbenfabriken Bayer A. G. Chemicals Corporate Licensee

DIRECTIONS FOR USE

Apply with low pressure sprayer equipped with a coarse spray nozzle. Spray until completely wet but not to runoff. Avoid excessive runoff. Apply to all insect breeding areas and harbors. Also treat adjacent exposed surfaces where insects are found when they come out of hiding.

REPEAT TREATMENT WHENEVER NECESSARY

ACTIVE INGREDIENTS

Methylene Chloride	1.00%
Petroleum Distillate	81.00%
INERT INGREDIENTS	15.00%

SEE SIDE PANEL FOR ADDITIONAL CONTENTS

DO NOT USE AS A SOLE SOURCE

CONTENTS: KEEP OUT OF REACH OF CHILDREN

(SEE SIDE PANEL FOR ADDITIONAL CONTENTS)

NET CONTENTS CAL.

BEST DOCUMENT AVAILABLE

Cockroaches, Crickets, Earwigs, Dog Fleas, Flies, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw Toothed Grain Beetles and Weevils (Exposed Stages) -

Silverfish, Firebrats and Pantries - Apply to all cracks and crevices inhabited by these insects.

Brown Dog Ticks - Spray sleeping quarters of pets, crevices and baseboards, wall and door frames, wall cracks, and floor areas of floors. Pay special attention to areas of pet sleeping quarters after spraying. Repeat spraying as needed.

Clover Mites, Crickets, Earwigs, Millipedes, Centipedes, Scorpions and Spiders - Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites, give special attention to windowsills.

Flies, Fleas, Gnats, Mosquitoes, Pantries and Silverfish (Outdoor Only) - Spray infested areas and the surrounding outside surfaces of the premises, including trees, fences, plants and other outdoor structures.

For Use in Food Processing and Food Preparation Areas - Use only in the food processing and food preparation areas. Apply only when food, food processing equipment, utensils, and food contact surfaces are not present or are covered before spraying. Remove all exposed food before reapplying on the food processing or food preparation areas. Apply only in the spaces between processing runs. Remove food and food contact surfaces before spraying. Do not spray food. Spray only the outside of food processing equipment. Spraying on loading docks and outside areas will help insect control and in preventing reinfestation. Good sanitation and proper disposal of waste is essential in an effective insect control program.

Insect Control in Research Facilities:

The use of Residual Spray in and around research animal facilities for insect control will not interfere with studies on Liver Microsomal Enzyme activity in laboratory animals.

10

16, 17

110