DIRECTIONS

Not For Use on Feed or Food Crops

WHEN TO APPLY: Applications can be made in all seasons. However, bait should not be applied in rainy periods, when rain is expected within 24 hours, or during periods of prolonged freezing weather.

HOW MUCH TO USE: Use at the rate of 1 pound for a medium-sized nest or "town" (one having a central nest area containing about 150 mounds). Use proportionate amounts for smaller or larger towns ranging from 6 pounds on very large, old "towns" to 1/4 pound on those having only a few mounds. For single-mound nests use 1 ounce

HOW TO APPLY: Scatter the pellets evenly over the central nest area where most of the excavation mounds are concentrated, and for the medium to large nests include an area about 20 feet wide bordering the central nest area. Bait can be broadcast with a suitable mechanical spreader or by hand. Wear rubber gloves when handling this product. Repeat applications as necessary.

HOW LONG FOR CONTROL: First signs of control will be a great reduction in excavation and foraging activities. This occurs in about 5 to 14 days. These activities gradually stop completely and the nest is dead in 5 to 6 weeks.

(Continued on Panel at right)

NET WEIGHT



The Te

CAUTIO For Additi

ACTIVE INGRED Dodecachloro pentalene* INERT INGREDI * A-C MIREX



MADE IN U. S. A.

SSACTA TRANSTERED From DIS (1)

States front	O POUNDS	(Continued from Panel at left) OTHER ANTS: This product w and FIRE ANTS in the treated a
		NOTE: Do not use other insect is being used; they may interfue bait.
LLETED BAIT	ACCEPT	ED CAU
``450''	MAR 6 1969	Keep out of r
For Control of	UNDER THE FEDERAL IN PE	Harmful if swallowed.
exas Leaf-Cutting A (TOWN ANT) M—Keep out of reach of ch tional Caution Information See Right Pane USDA Reg. No. 218-	33462-9 ildren	Wash with soap and we eating or smoking. Do not contaminate feed (Keep out of reach of chil
DIENT poctahydro-1,3,4-metheno-2H-cyclobuta[cd] IENTS		Wear rubber gloves wher Keep Container Tightly C Dry Place Away From OI
U.S. Pet. Reissue 24,750		DO NOT REUSE
ALLIED CHEMICAL CORPORATION DEPARTMENT AG 40 RECTOR STREET, NEW YORK, N.Y., U.S.A.		Follow directions carefully. Timin and crop conditions, mixtures w recommended, and other influenci are beyond the control of the se storage or handling of this materi
9481 A 15088 PF	INTED IN U. S. A.	tions given herewith.

9481 A 1501 8

PRINTED IN U. S. A

S

Food Crops


be made in all seasons. rainy periods, when rain r periods of prolonged

rate of 1 pound for a ing a central nest area portionate amounts for 5 pounds on very large, ing only a few mounds.

evenly over the central on mounds are concennests include an area al nest area. Bait can be preader or by hand. Wear uct. Repeat applications

signs of control will be foraging activities. This activities gradually stop to 6 weeks.

Continued on Panel at right)



MADE IN U. S. A.

A 1501 B

PRINTED IN U. S. A.

ELKELD TROM 218-623

(Continued from Panel at left)

OTHER ANTS: This product will also control HARVESTER ANTS and FIRE ANTS in the trested area.

NOTE: Do not use other insecticides on the area while the bait is being used; they may interfere with the effectiveness of the bait

CAUTION

Keep out of reach of children

Harmful if swallowed.

FOR ECONOMIC PUSON REGISTER. I

Wash with soap and water after using and before eating or smoking.

Do not contaminate feed or foodstuffs.

Keep out of reach of children and domestic animals.

Wear rubber gloves when handling.

Keep Container Tightly Closed And Stored In a Cool Dry Place Away From Other Pesticides.

DO NOT REUSE THIS CONTAINER

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.