

Master Label

Ready-to-Use

Y-TEX®

BRUTE™

Pour-on for Cattle and Swine

- For long-lasting residual control of horn flies, lice and Gulf Coast ticks on lactating and non-lactating dairy cattle and beef cattle.
- For control of lice and mange mites on swine.
- Precise dose dispensed directly from container or dispensing gun.
- Concentrate for dilution with diesel fuel or oil for backrubbers.
- Recommended for Agricultural/Commercial use.

Active Ingredient

Permethrin* 10.0%
 (3-Phenoxyphenyl)methyl(±)cis-trans 3-(2,2-dichloroethyl)-
 -2,2-dimethylcyclopropanecarboxylate.

INERT INGREDIENTS† 90.0%
 100.0%

*cis/trans ratio: Min. 35%(±)cis and Max. 65%(±) trans.

†Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE INSIDE OF BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS:

EPA Est. No. 10370-TX-1 EPA Reg. No. 39039-7

Y-TEX CORPORATION • 1825 Big Horn Avenue • Cody • WY 82414

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

Harmful if inhaled. Causes moderate eye injury and skin irritation. Do not get on skin or clothing. Avoid breathing vapor or spray mist and contact with eyes. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Do not use on horses.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Promptly drink a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately. Do not induce vomiting unless under a physician's care. This product contains petroleum distillate. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove victim to fresh air, apply artificial respiration if indicated. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

NOTICE OF WARRANTY:

Y-TEX CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label, and in any case, is limited to replacement of product only.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL & CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES:

Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL:

Containers one gallon and smaller: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Containers larger than one gallon: Metal containers – Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers – Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities; by burning, if burned, stay out of smoke. Glass containers – Triple rinse or equivalent. Then dispose of in a sanitary landfill or by other approved state and local procedures.

ACCEPTED

FEB 25 1998


Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 39039-7

BACK PANEL LABEL TEXT SPECIFIC TO CONTAINER
WITH MEASURING CHAMBER INCORPORATED INTO CONTAINER:

- Ready-to-use pour-on to control horn flies, lice and Gulf Coast ticks on cattle.
- Ready-to-use pour-on to control lice and mange mites on swine.
- Concentrate for dilution with diesel fuel or oil for backrubbers.
- Can be used on lactating dairy cattle with no withholding period.
- As a pour-on, one pint treats:

WEIGHT (lbs.)	No. of CATTLE	No. of SWINE
200	157	78
400	78	40
600	52	32
800	40	-
1000	32	-

Store upright at room temperature. Avoid exposure to extreme temperatures. Do not contaminate water, food or feed by storage. See inside for complete Storage and Disposal instructions.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE ON CATTLE

Apply to Dairy Cattle (lactating and non-lactating) and Beef Cattle. Shake well before using. Loosen cap, squeeze bottle to fill measuring chamber with required dosage. Remove cap and apply to animal.

TARGET INSECTS	APPLICATION INSTRUCTIONS
For up to 6 weeks' control of susceptible horn flies.	Apply down midline of back over shoulders.
For control of lice, Gulf Coast ticks and aid in control of other tick species, face flies, stable flies, horse flies, deer flies, house flies, mosquitoes and black flies.	Apply from poll down neck, over shoulders. Dose: (ml or cc) Weight: (lb.)
	3 200
	6 400
	9 600
	12 800
	15 1000 & over

Repeat treatment as needed, but not more often than once every 2 weeks. For optimum lice control, two treatments at a 14 day interval are recommended. Do not use on horses. Special Note: Brute is non-systemic and will not control cattle grubs.

DIRECTIONS FOR USE ON CATTLE CONTINUED

TARGET INSECTS	APPLICATION INSTRUCTIONS
For control of horn flies, lice, face flies, and stable flies.	Backrubber: Dilute 48 ml (1.6 fl. oz.) in 1 gal. diesel fuel or oil. Keep rubbing device charged. Results can be improved by forced use.

DIRECTIONS FOR USE ON SWINE

Apply to Swine (100 lbs. and over). Shake well before using. Loosen cap, squeeze bottle to fill measuring chamber with required dosage. Remove cap and apply to animal.

TARGET INSECTS	APPLICATION INSTRUCTIONS
For control of lice, mange mites and aids in control of biting and nuisance flies.	Apply across back of head and ears, then apply down midline of neck, over shoulders. Dose: (ml or cc) Weight: (lb.)
	3 100
	6 200
	9 300
	12 400
	15 500 & over

Repeat treatment as needed, but not more often than once every 2 weeks. For optimum control, two treatments at a 14 day interval are recommended. Allow 5 days between last treatment and slaughter.

BACK PANEL LABEL TEXT SPECIFIC TO CONTAINERS
WITH A SEPARATE APPLICATOR GUN:

- Ready-to-use pour-on to control horn flies, lice and Gulf Coast ticks on cattle.
- Ready-to-use pour-on to control lice and mange mites on swine.
- Concentrate for dilution with diesel fuel or oil for backrubbers.
- Can be used on lactating dairy cattle with no withholding period.
- As a pour-on, one gallon treats:

WEIGHT (lbs.)	No. of CATTLE	No. of SWINE
199	1,514	757
399	757	378
599	504	252
799	378	-
1000	252	-

Store upright at room temperature. Avoid exposure to extreme temperatures. Do not contaminate water, food or feed by storage. See inside for complete Storage and Disposal instructions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE ON CATTLE

Apply to Dairy Cattle (lactating and non-lactating) and Beef Cattle. Shake well before using.

TARGET INSECTS	APPLICATION
For up to 6 weeks' control of susceptible horn flies.	Apply down midline of back over shoulders.
For control of lice, Gulf Coast ticks and aid in control of other tick species, face flies, stable flies, horse flies, deer flies, house flies, mosquitoes and black flies.	Apply from poll down neck, over shoulders. Dose: (ml or cc) Weight: (lb.)
	2.5 Birth-199
	5.0 200-399
	7.5 400-599
	10.0 600-799
	12.5 800-999
	15.0 1000 & over

Repeat treatment as needed, but not more often than once every 2 weeks. For optimum lice control, two treatments at a 14 day interval are recommended. Do not use on horses. Special Note: Brute is non-systemic and will not control cattle grubs.

For control of horn flies, lice, face flies, and stable flies.	Backrubber: Dilute 48 ml (1.6 fl. oz.) in 1 gal. diesel fuel or oil. Keep rubbing device charged. Results can be improved by forced use.
--	--

DIRECTIONS FOR USE ON SWINE

Apply to Swine (85 lbs. and over). Shake well before using.

TARGET INSECTS	APPLICATION
For control of lice, mange mites and aids in control of biting and nuisance flies.	Apply across back of head and ears, then apply down midline of neck, over shoulders. Dose: (ml or cc) Weight: (lb.)
	2.5 85-169
	5.0 170-254
	7.5 255-339
	10.0 340-424
	12.5 425-509
	15.0 510 & over

Repeat treatment as needed, but not more often than once every 2 weeks. For optimum control, two treatments at a 14 day interval are recommended. Allow 5 days between last treatment and slaughter.

