


# Y-TEX CORPORATION

P. O. Box 1450, 1825 Big Horn Avenue, Cody, Wyoming 82414-1450 • Phone (307) 587-5515 • FAX (307) 527-6433



## TUTTLE'S® **BRUTE**®

Insecticide for Horses

- Long-Lasting Control of Biting Flies, Including Mosquitoes, on Horses.
- Reduces Bot Fly Eggs Laid on Horses for up to 4 Days.
- Kills and Repels Black Flies and Ticks.
- Precise Dose Directly from Container.
- Use In and Around Livestock Premises as a Spot Treatment.

EPA Est. No. 39578-TX-01 EPA Reg. No. 39039-7  
TUTTLE'S HORSE CARE DIVISION,  
Y-TEX Corporation, 1825 Big Horn Ave., Cody, WY 82414



**ACCEPTED**

JUN - 1 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 39039-7

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
& DOMESTIC ANIMALS  
WARNING**

Harmful if inhaled. Causes moderate eye injury and skin irritation. Do not get on skin or clothing. Avoid breathing vapor or spray mist and contact with eyes. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

**STATEMENT OF PRACTICAL TREATMENT**  
**IF SWALLOWED:** Promptly drink a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately. Do not induce vomiting unless under a physician's care. This product contains petroleum distillate. Vomiting may cause aspiration pneumonia.

**IF INHALED:** Remove victim to fresh air, apply artificial respiration if indicated. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF IN EYES:** Flush with plenty of water. Call a physician if irritation persists.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to fish. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**PHYSICAL & CHEMICAL  
HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**NOTICE OF WARRANTY:**

Y-TEX CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label, and in any case, is limited to replacement of product only.

YTP 125 400

**TUTTLE'S®  
BRUTE®  
Insecticide for Horses**

- Long-Lasting Control of Biting Flies, Including Mosquitoes, on Horses.
- Reduces Bot Fly Eggs Laid on Horses for up to 4 Days.
- Kills and Repels Black Flies and Ticks.
- Precise Dose Directly from Container.
- Use in and Around Livestock Premises as a Spot Treatment.

EPA Est. No. 39578-TX-01 EPA Reg. No. 39039-7  
TUTTLE'S HORSE CARE DIVISION,  
Y-TEX Corporation, 1825 Big Horn Ave., Cody, WY 82414



3/8

### DIRECTIONS FOR USE ON HORSES

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For dirty horses, shampoo and rinse thoroughly. Wear protective gloves or protective application mitt when using as a wipe-on for horses. Shake well before using. Loosen cap, squeeze bottle to fill measuring chamber with required dosage. Remove cap and apply to animal. **Apply 3-6 ml per 200 lb. body wt. up to a max. of 30 ml. Do not treat foals under 12 weeks of age.**

APPLICATION METHOD	TARGET PEST	APPLICATION INSTRUCTIONS
Ready-to-Use Wipe-on (Preferred method for optimum control of most pest species).	For control of mosquitoes, black flies, face flies, horn flies, lice and ticks. Aids in control of stable flies, horse flies, deer flies, house flies, bot flies and eye gnats.	Apply as a whole body wipe with cloth or insecticide application mitt. For the first application with a new cloth or mitt apply twice the dosage to charge the application cloth or mitt. Pay special attention to where pests congregate. • Face flies and eye gnats - apply to face and around eyes being careful not to get product in the eyes. • Black flies - wipe the inside of the ears, under chin, throat and midline of belly. • Stable flies - wipe legs, lower shoulders and belly. • Bot flies - wipe under chin, throat, chest and legs.
Ready-to-Use Pour-on.	For control of horn flies.	Apply over withers and down back. Do not apply to horses that will be ridden within 24 hours of application. Pour-on application may leave an oily residue on the coat of some animals.

Repeat treatment as needed but not more often than once every 7 days.

### DIRECTIONS FOR USE IN AND AROUND LIVESTOCK PREMISES

For use in and around livestock premises as a spot treatment. Shake well before using. Loosen cap, squeeze bottle to fill measuring chamber with required dosage. Remove cap and apply.

TARGET INSECTS	APPLICATION INSTRUCTIONS
For control of house flies, stable flies and other nuisance pests in and around stables and other livestock premises.	Apply undiluted as a coarse spray or wipe directly to surface where pests crawl and alight. One fl. oz. will treat 560 ft <sup>2</sup> . One pint will treat 8900 ft <sup>2</sup> .

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage and Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**Pesticide Disposal:** Pesticide that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**Container Disposal:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

3/3