

BALCOM



REGISTERED

SEP 8 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 960-166

changed to:
34704-7

BUTYL 6 Ester Weed Killer

EMULSIFIABLE CONCENTRATE FOR CONTROLLING BROAD-
LEAF WEEDS IN CORN, WHEAT, BARLEY, RYE, AND NON-CROP
AREAS.

CAUTION: Keep Out Of Reach Of Children.

CAUTION: MAY CAUSE SKIN IRRITATION.

AVOID CONTACT WITH EYES, SKIN, OR CLOTHING. HARMFUL
IF SWALLOWED.

- Do not store near food, fertilizers, seeds, insecticides, or fungicides.
- Do not contaminate water used for irrigation, domestic or spray purposes.
- This product is toxic to fish. Keep out of lakes, streams or ponds.
- Do not contaminate water by cleaning of equipment, or disposal of wastes.

ACTIVE INGREDIENTS:

*Isobutyl ester of	
2,4-Dichlorophenoxyacetic acid	51.4%
*n-Butyl ester of	
2,4-Dichlorophenoxyacetic acid	26.5%

INERT INGREDIENTS:	22.1%
	100.0%

*Equivalent to 62.2% 2,4-D acid or 6.0 pounds per gallon.
Store above 5° F. to keep from freezing.
If allowed to freeze, warm to 70° F. and dissolve before using.

NET CONTENTS U.S. GALLON(S).

USDA Reg. No. 960 -166

DIRECTIONS FOR USE

Because of variations in climate or other factors, there are various expert opinions as to the best rate of application. Consult experiment station or Extension Service weed specialists for their specific recommendations on type of application, dosages, and timing of use. You may wish to follow their recommendations, which may vary somewhat from those given below.

CAUTION: DO NOT USE UNTIL YOU HAVE READ THE "WARNING" ON THIS LABEL.

WHEAT, BARLEY, OR RYE—To control easy-to-kill weeds, use 1/3 pint per acre. To control hard-to-kill weeds, apply 1/2 pint per acre. Treat after tillering begins and before crop emerges from the boot or before jointing. Treat winter wheat only in spring when weeds are small, and before the crops begin heading. Application at other times may cause abnormal heading. Do not spray these crops when seeded down with legumes. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.

CORN—Apply 1/3 pint per acre. Treat any time before tassels appear, but if corn is 8 inches or taller, avoid application to the growing point by using drop nozzles and directing the spray near the base of the crop. Because different varieties of corn vary in susceptibility to herbicides, consult competent local agricultural authorities to determine if use is safe in your particular hybrid.

Avoid cultivating soon after treatment when corn may be brittle. Do not use in corn grown for seed production. For pre-emergence treatment, use 2 2/3 pints overall per acre; spray one to three days before corn emerges. Do not use pre-emergence sprays on light, sandy soils.

LAWNS, GOLF COURSES AND SIMILAR TURF—Apply 1 1/3 pints per acre. For small areas, use 2 teaspoonfuls per gallon of water. Do not apply to newly seeded lawns until grass is well established. Do not use on bent or other creeping grasses. Clover or other legumes may be injured by spraying, especially during hot summer weather. Avoid spraying seed grasses when in the blossoming or pollinating stage. Spray or vapors from this product may injure Dichondra.

NON-CROP AREAS (FENCEROWS, DITCHBANKS, RIGHTS-OF-WAY AND SIMILAR AREAS)—Apply 1 1/3 to 2 pints in 100 gallons of water and wet foliage thoroughly. Repeat treatments may be needed to control hard-to-kill weeds; re-treat when regrowth is visible.

WOODY PLANTS—To control susceptible woody plants like Willow, Sumac, Buckbrush, use 2 1/2 to 4 pints per hundred gallons of water and wet all foliage thoroughly. Treat when brush is growing rapidly, in late spring or early summer. For small areas, use 4 teaspoonfuls per gallon of water.

To prevent cut stumps from resprouting, mix one gallon in 25 to 30 gallons of No. 2 fuel oil or diesel oil. Wet stump thoroughly down to ground line. For treating small areas, mix 1/4 pint to 1 gallon of oil. Stump treatments may be made at any time of year.

SPOT TREATMENT: For spot treatment spraying of small non-crop areas, mix 1 tablespoon per gallon of water and thoroughly wet all weed foliage.

WARNING NOTICE:

Balcom Chemicals, Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Balcom Chemicals, Inc. makes no other express or implied warranty of fitness or merchantability. The exclusive remedy of the user or buyer, and the limit of the liability of Balcom Chemicals, Inc. or any other seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer of the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice, which may not be varied by any verbal or written agreement.

34704-1

BALCOM CHEMICALS, INC.

