



Net Weight 50 Pounds

PREMIUM  
SUPER PIG STARTER 500 (ROL)  
Contains Rabon<sup>®</sup> Oral Larvicide

To prevent the development of house flies in the manure of treated Swine.

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate <sup>®</sup>	0.02%
INERT INGREDIENTS <sup>**</sup>	99.98%
TOTAL	100.00%

<sup>®</sup>Rabon Insecticide. Contains 90.8 mg. Rabon per pound.  
<sup>\*\*</sup>Refers only to ingredients which are not larvicidal.  
Rabon is a registered trademark of Shell Oil Company.

GUARANTEED ANALYSIS

Crude Protein. . . . .	.Min. 15.00%
Crude Fat. . . . .	.Min. 2.00%
Crude Fiber. . . . .	.Max. 2.00%

FEED INGREDIENTS

Animal Protein Products, Process Grain By-Products, Potassium Chloride, Dicalcium Phosphate, Calcium Carbonate, Magnesium Oxide, Salt, Bentonite, DL-Methionine, L-Lysine, Sorbic Acid, Natural and Artificial Flavors, Vitamin A Acetate in Gelatin, Vitamin D3 Supplement, Vitamin E Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K), Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Vitamin B12 Supplement, Folic Acid, d-Biotin, Ascorbic Acid, Choline Chloride, Zinc Oxide, Ferrous Sulfate, Copper Sulfate, Manganous Oxide, Copper Oxide, Ethylenediamine Dihydrochloride, Cobalt Carbonate, and Sodium Selenite.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place.

PESTICIDE DISPOSAL: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if

allowed by state and local authorities, by burning. If burned, stay out of

smoke

MANUFACTURED BY:  
CARL S. AKEY, INC.  
LEWISBURG, OHIO 45338

EPA Reg. No. 39258-

EPA Est. No. 39258-OH-1

39258-4

PREMIUM  
SUPER PIG STARTER 500 (ROL)

DIRECTIONS FOR USE  
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Start feeding RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of house flies in treated swine, but it is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around barns to control adult house flies which can breed not only in manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, this product should be used in conjunction with other good management and sanitation practices.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix.

Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixer, horizontal blenders,) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

MIXING AND FEEDING DIRECTIONS

PIGS: For starting pigs mix 500 lbs. of this premix with 1,150 lbs. of ground corn and 350 lbs. of soybean meal containing 44% protein. Feed the resulting larvicidal feed as the sole ration. This feed is designed as a pre-starter and starter feed containing approximately 17% protein and 22.7% of Rabon per lb. All swine should be treated for best results.

<p>U.S. ENVIRONMENTAL PROTECTION AGENCY</p> <p>NOTICE OF PESTICIDE</p>	<p>JUN</p>
<p>NAME AND ADDRESS OF REGISTRANT</p> <p style="text-align: right;">160 / 117644 18 / 1</p>	
<p><b>NOTE</b> Changes in labeling formula, formula, or substance that are accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>	
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p>	
<p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p>	
<p>Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his own motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>	
<p><input type="checkbox"/> ATTACHMENT SHEET(S) ATTACHED</p>	
<p>SIGNATURE OF APPLICANT</p>	<p>DATE</p>