Net Weight 50 Pounds

PREMIUM SWINE 50 (BOL) PREMIX Contains Rabon Oral Larvicide

To prevent the development of house flies in the manure of treated Swine.

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate*	0.20%
INERT INGREDIENTS**	99.80%
TOTAL.	100.00%

*Rabon Insecticide. Contains 0.9 grams Rabon per pound. **Refers only to ingredients which are not larvicidal. Rabon is a registered trademark of Shell Oil Company.

GUARANTEED ANALYSIS

Calcium (Ca) .	٠	٠	•	٠	٠	•	•	•	•	•	٠	•	٠	•	•	.Min.	19.0%	Max.	22.8%
Phosphorus (P)	•	•	•	•	•	•	٠	•	•	•	•	•	•	•	٠	.Min.	6.9%		
Salt (NaCl)	•	•	•	•	•	•	•	٠	•	٠	•	•	•	٠	•	.Min.	13.0\$	Max.	15.6%

FEED INGREDIENTS

Dicalcium Phosphate, Calcium Carbonate, Salt, Potassium Chloride, Magnesium Oxide, Vitamin A Acetate in Gelatin, Vitamin D3 Supplement, Vitamin E Supplement, Menadione Dimethylpyrimidinol Bisulfite, (Source of Vitamin K), Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Vitamin B12 Supplement, Zinc Oxide, Ferrous Sulfate, Manganous Oxide, Copper Oxide, Ethylenediamine Dihydriodide, Sodium Selenite, and Cobalt Carbonate.

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, see a physician.

ENVIRNOMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. <u>STORAGE</u>: Store in a dry place.

<u>PESTICIDE DISPOSAL</u>: Wastte resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

<u>CONTAINER DISPOSAL</u>: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, (f allowed by state and local authorities, by burning. If burned, stay out of smoke.

 $\frac{ACCOPTED}{w = 0 + 200 \text{ NES}}$ in LEV 1 = 0.1 \pm 1

MANUFACTURED BY: CARL S. AKEY, INC. LEWIGBURG, OHIO 45338

JUN 1 *224

EPA Reg. No. 39258-

EPA Est. No. 39258-0H-1

251012 . 1.

PREMIUM SWINE 50 (ROL) PREMIX

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Start feeding RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of house flies in treated swine, but it is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around barns to control adult house flies which can breed not only in manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, this product should be used in conjunction with other good management and sanitation practices.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix.

Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixer, horizontal blenders,) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

MIXING AND FEEDING DIRECTIONS

SWINE (125 lbs. - market weight): Mix 50 lbs. of this premix with 1950 lbs. of ground corn and soybean meal as in the chart below. The resulting larvicidal feed contains 22.7 mg. of Rabon per pc ind. Feed as the sole ration. This ration may be fed up to slaughter. All swine should be treated for best results.

Protein % Hog Wt., Lbs.	1 125-	4 •175	13 175-Mkt.		
Ground Corn	1650	1680	1700	1730	
SBM (44%)	300		250		
SBM (43 \$)		270		220	
Swine 50 (ROL) Premix	50	50	50	50	
Total Lbs.				2000	

THAT ON NO PAREGE P A U.S. ENVIRONMENTAL PROTECT DN GENCY OFFILE OFFE TEPM OF FSUANCE NAME OF PETTICLE PRODUCT NOTICE OF PESTICIDE ۰. NAME AND ADDRESS OF RE A THAN? In Frank 210 - 1 Г L .1 NOTE Changes in labeling formula differing in substance list, that accepted in connection with this registration must be submitted to and accepted by the Registration Division grants user of the label in coemerce. In any correspondence on this product always refer to the above U.S. E.P.A reportation numbers. On the Lass of information furnish, dity the registrant, the above named posticide is hereby Registered/Reregistered under the Federal Intesticate, $1~{\rm mpc}$ ade, and R shouts als A(t)A copy of the labeling accepted in connection with this Keyle tration Receptionation is returned herewith. Registration is in no way to be construct as an inderservent or approval of this product by this Agency. In order to protect health and the environment, the A ferencipation on his postion, carved any time suspend or cancel the registration of a pestscide in accordance with the A-t. The a section vision matching encoder in with the registration of a product under this Act as not to be a true to pay ty the reputrient a tradit several vessel of the name of to its use if it has been covered by others. , SPOT DOOLSHIT IN HAPS ATTA HMEN'S ANNU ENLY DATE SIGNAT GETEADDUT, N. F. A. .

• ··· ·· ··