Net Weight 50 Pounds

PREMIUM SOW - 100 (BOL) PREMIX Contains Rabon Oral Larvicide

To prevent the development of house flies in the manure of treated Swine.

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate*	0.20%
INERT INGREDIENTS**	99.80%
TOTAL	100.00%

*Rabon Insecticide. Contains 0.9 grams Rabon per pound.

**Refers only to ingredients which are not larvicidal.

Rabon is a registered trademark of Shell Oil Company.

GUARANTEED ANALYSIS

Calcium (Ca)	 	16.00% Max. 19.00%
Phosphorus (P)	 	9,50%
Salt (NaCl)	 	8.40% Max. 10.00%

FEED INGREDIENTS

Dehydrated Alfalfa Meal, Potassium Chloride, Dicalcium Phosphate, Calcium Carbonate, Salt, Potassium Sulfate, Magnesium Sulfate, Vitamin A Acetate in Gelatin, Vitamin D₃ Supplement, Vitamin E Supplement, Menadione Dimethylpyrimidinol Brsulfite (Source of Vitamin K), Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Vitamin B₁₂ Supplement, Folic Acid, d-Biotin, Choline Chloride, Zinc Oxide, Ferrous Sulfate, Manganous Oxide, Copper Oxide, Ethylenediamine Dihydriodide, Cobalt Carbonate and Sodium Selenite.

with COMMENTS In EPA Letter Date di

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS
CAUTION

JUN 1 1984

Urder the in

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before that or smoking. If in eyes, wash with plenty of water. If irritation 3935 persists, see a physician.

ENVIRNOMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a mrry place.

PESTICIDE DISPOSAL: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration cr, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

MANUFACTURED BY: CARL S. AKEY, INC. LEWISBURG, OHIO 45338

EPA Reg. No. 39253-

EPA Est. No. 39258-0H-1

PREMIUM SOW - 100 (ROL) PREMIX 3100 51

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Start feeding RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

When fed, this product passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch.

It prevents the development of house flies in treated swine, but it is not effective against existing adult flies.

In some cases, supplemental fly control measures may be needed in and around barns to control adult house flies which can breed not only in manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, this product should be used in conjunction with other good management and sanitation practices.

This product will mix uniformly in feeds when standard mixing procedures are followed. Thus, usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution throughout the feed mix.

Optimum performance can be assured only if this product is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixer, horizontal blenders,) may be used to prepare formulated feeds.

Feeds prepared with this product should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, this product should not be mixed in liquid feed supplements.

MIXING AND FEEDING DIRECTIONS FOR SOWS

Mix 100 lbs. of this product with 1900 lbs. of ground corn and soybean meal using formulation A or B. The resulting larvicidal feed contains approximately 14.3% protein and 45.4 mg. of Rabon per pound. Offer 4 to 6 pounds per sow daily. This product may be fed up to slaughter.

INGREDIENT	A	B
Ground Corn (8.9% Protein)	1575 lbs.	1600 lbs.
Soybean Meal (44% Protein)	325 lbs.	
Soybean Meal (48.5% Protein)		300 lbs.
Premium Sow 100 (ROL) Premix	100 lbs.	100 lbs.
TOTAL MIX	2000 lbs.	2000 lbs.

	J+P 5/29/g
U.S. ENVIRONMENTAL PROTECTION AGENCY OFF OF THE THE PROTECTION ATT OUR THAT TOWN ENVIRONMENTAL ATT OUR THAT TOWN ENVIRONMENTAL ATT	FERM OF I SUANCE
NOTICE OF PESTICIDE: $\frac{n_{t}}{n_{t}} = \frac{n_{t}}{n_{t}} = \frac{n_{t}$	MEANING OF THE FORMS DON'T
NAME AND ADDRESS OF REGION AND THE THE STIP TOLEGO	<u> </u>
r- ,	18/1
L.	.1
	from that accepted in connection with this registration must be a to use of the latel in commer e. In any correspondence on this for.
On the basis of information farmished by the registrant, the the Federal Insection is, I uncould, and R shown ide Act.	al we named posticide is hereby Registered/Reregistered under
A copy of the labeling accepted in connection with this Re-	gestrate in Receptoration is returned betewith.
health and the environment, the Alb involvator in his motivicide in accordance with the Alb. The accorptance of an aba	it or approval of this product by this Ayency. In order to protect in, may it any time suspend or cancel the registration of a pestame in innection with the registration of a product under this leastly use to of the name or to its use if it has been covered.
	·
ATTA NUMBER AND AND	
SIGNATURE FADDE LIGHT E.	DAYE

BEST DOCUMENT AVAILABLE