

Net Contents 30 Gallons/113.6 Liters



Diamond Shamrock

ACCEPTED
OCT 13 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act
39335-43

2,4-D Butyric

Weed Killer
Herbicide

Active Ingredient: Dimethylamine salt of 4-(2,4-dichlorophenoxy) butyric acid*	25.9%
Inert Ingredients:	74.1%
Total:	100.0%

*4-(2,4-dichlorophenoxy) butyric acid equivalent to 21.9%; 2 lbs/gal.

Keep Out of Reach of Children

CAUTION

See side panel for additional precautionary statements

Diamond Shamrock Agricultural Chemicals, Inc.

P.O. Box H
Tuscaloosa, AL 35404

EPA Reg. No. 39335-43

EPA Est. No. 39335-AL-1



39335-43

10/13/81

1/2

Precautionary Statements

Hazards to Humans and Domestic Animals

Caution

Harmful if swallowed or absorbed through the skin. If swallowed, **Do not** induce vomiting. If vomiting occurs spontaneously, keep airway clear. Never give anything by mouth to an unconscious person. Seek medical attention.

Causes eye and skin irritation.

Do not inhale spray mist. **Do not** get in eyes. In case of contact flush eyes with plenty of water for at least 15 minutes and seek medical attention. **Do not** allow contact with skin or clothing. Wash exposed skin gently with plenty of soap and water.

Environmental Hazards

Do not apply directly to water. **Do not** contaminate water by cleaning of equipment or disposal of wastes. Use care to avoid spray contact or drift to susceptible plants such as beans and other legumes, cotton, flowers, grapes, ornamentals, vegetables and other plants. **Do not** permit spray mist containing 2,4-D Butyric Weed Killer to drift onto them, since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. **Do not** spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 8 miles per hour. **Do not** apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray. With aircraft application, drift can be lessened by applying not less than 5 gallons of spray per acre; by using no more than 20 pounds spray pressure at the nozzles; by using nozzles which produce a coarse spray pattern; and by spraying only when the wind velocity is less than 5 miles per hour. Flush sprayer out on suitable non-crop area after use. **Do not** use the same spray equipment for applying other materials to susceptible crops as injury may result.

2,4-D Butyric Weed Killer

Read entire label before using this product.

General Information

2,4-D Butyric Weed Killer is recommended for the control of many broadleaved weeds in seedling or established legumes. The following is a partial list of weeds controlled by 2,4-D Butyric Weed Killer:

Canada thistle	Nightshade
Cocklebur	Pigweed
Curly dock (seedling)	Ragweed
Field bindweed	Russian thistle
Kochia	Smartweed
Lambsquarter	Stinkweed (pennycress)
Morningglory	Whitetop (hoary cress)

Note: Local conditions, crop varieties and application regulations vary and may affect use of this herbicide. Consult local agricultural experiment station or extension service weed specialists and state regulatory agencies for recommendations in your area.

Note: When stored at temperatures below freezing, it may be necessary to warm contents to 45°F and mix thoroughly before using.

NOTE: DO NOT graze established crop, or feed straw or hay from treated crop to livestock within 30 days after application. DO NOT graze or feed hay from treated seedling crops to livestock within 60 days after application.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Preparation of the Spray: Fill the spray tank with half the required amount of water, then add the recommended amount of 2,4-D Butyric Weed Killer with agitation and continue filling the spray tank with water. Use 5 to 10 gallons of water per acre or sufficient water to give uniform coverage.

Time of Application: Application should be made on seedling or established legumes such as alfalfa, redclover, trefoil and white clover in Spring or Fall 2 to 4 weeks after emergence of legumes, and when seedling weeds are less than 1 to 2 inches high or have rosettes less than 3 inches across. Later applications give unsatisfactory control.

Apply 2,4-D Butyric Weed Killer 1 to 2 quarts per acre to annual weeds less than 1 inch tall and 2 to 3 quarts per acre to annual weeds 1 to 3 inches tall. For control of perennial weeds up to 3 inches tall apply 2 to 3 quarts per acre.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal: **Do not** reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved State and local procedures.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. **Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty.** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.