

Sumithion® 40WE

FOR USE ONLY BY OR UNDER THE SUPERVISION OF STATE OR FEDERAL OFFICIALS RESPONSIBLE FOR MALARIA CONTROL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes eye irritation. Harmful if swallowed. Do not get in eyes. Avoid contact with skin. Wear protective gloves, clean protective clothing and goggles or face shield when handling the material. Remove and wash contaminated clothing before reuse. Wash hands and arms thoroughly after handling and before eating, drinking or smoking.

STATEMENT OF PRACTICAL TREATMENT

If in eyes, wash thoroughly with water for 15 minutes. Get medical attention.

If contact with skin, wash thoroughly with soap and water. Get medical attention if irritation persists.

Note to Physician: Sumithion is an organophosphorus compound that inhibits cholinesterase. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Do not permit children or pets near treated surfaces until dry.

Keep out of domestic or animal water supplies.

Do not store or transport with food or animal feed.

Do not contaminate water by cleaning equipment or disposal of containers.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.
- 2. PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- 3. CONTAINER DISPOSAL:** Dispose of in a incinerator or landfill approved for pesticide containers, or bury in a safe place.
- 4. GENERAL:** Consult Federal, State or local disposal authorities for approved alternative procedures such as incineration or open burning.

ACTIVE INGREDIENTS

O,O-Dimethyl O-4-nitrophenyl phosphorothioate

INERT INGREDIENTS

40% (w/w)

60% (w/w)

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

If in eyes, wash thoroughly with water for 15 minutes. Get medical attention.

If contact with skin, wash thoroughly with soap and water. Get medical attention if irritation persists.

Note to Physician: Sumithion is an organophosphorus compound that inhibits cholinesterase. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DIRECTIONS FOR

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For control of adult Anopheles only. Not for use in food handling areas of private residences.

Apply as a residual spray. 1 gram (active ingredient) (0.1 floor area, furniture and other is complete, remove cover landfill approved for pesticide occupants of residence to.

Recommended dosage: Mix with 8 liters (2.1 gal) of water surface to obtain 2 g/sq m. Do not apply more often than

NOTICE OF CONDITIONS OF SALE

Sumitomo (and seller) offer(s) to users are deemed to have accepted warranty which may only be varied by representative of Sumitomo.

WARRANTY LIMITATION

Sumitomo warrants that this product meets the directions for use on the label. Sumitomo makes no other WARRANTY OF MERCHANTABILITY beyond the description on the label.

INHERENT RISKS:

The directions for use of this product should be followed carefully. However, with use. Buyer assumes all product contrary to label instructions conditions.

LIMITATION OF LIABILITY

In no case shall Sumitomo be liable for damages resulting from the use or handling of this product in an amount greater than the purchase price of such damages are claimed.

EPA Registration No. 147-000000
EPA EST. No. 147-000000

76

SUMITOMO CHEMICAL AMERICA, Inc.

343 PARK AVENUE
NEW YORK, N.Y. 10154

DO NOT USE IN RESIDENCES

Sumithion^(R) 40WDP

USE ONLY BY OR UNDER THE SUPERVISION OF STATE OR FEDERAL OFFICIALS RESPONSIBLE FOR MALARIA CONTROL

AC 1000000
1000000
1000000

PESTIC ANIMALS

not get in eyes. Avoid
n protective clothing
material. Remove and
wash hands and arms
drinking or smoking
TMENT
minutes. Get medical
water. Get medical

corus compound that
PAM is also antidotal
tropicine
faces until dry

ed
ent or disposal of

water, food or feed
bited.

spray mixture or
reprocessed. Should
indies or buried in a

or in a incinerator
bury in a safe place
or local disposal
ies, such as, etc.

ACTIVE INGREDIENTS

O,O-Dimethyl O-(4-nitro-m-tolyl)
phosphorothioate

INERT INGREDIENTS:

40% (w/w)
60% (w/w)

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR PRECAUTIONARY STATEMENTS

STATEMENT OF PRACTICAL TREATMENT

If in eyes, wash thoroughly with water for 15 minutes. Get medical attention.

If contact with skin, wash thoroughly with soap and water. Get medical attention if irritation persists.

Note to Physician: Sumithion is an organophosphorus compound that inhibits cholinesterase. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner not consistent with its labeling.

For control of adult Anopheles mosquitoes in private residence only. Not for use in food handling establishments or in food handling areas of private residences.

Apply as a residual spray to wall surfaces at a rate not to exceed 2 grams (active ingredient) (0.07 oz) per square meter of surface. Cover floor area, furniture and other objects prior to spraying. After spraying is complete, remove coverings and dispose of in an incinerator or landfill approved for pesticides or bury in a safe place. Do not allow occupants of residence to return until treated surfaces have dried.

Recommended dosage: Mix one kilo (2.2 lbs) of Sumithion 40 WDP with 8 liters (2.1 gal) of water and apply to 200 sq m (2153 sq ft) of wall surface to obtain 2 g/sq m (0.07 oz/10.8 sq ft). Agitate well before using. Do not apply more often than once every three months.

NOTICE - READ CAREFULLY

CONDITIONS OF SALE:

Sumitomo and seller warrant this product for sale subject to, and buyer and user are deemed to have accepted, the following conditions of sale and warranty, which may only be varied by written agreement of a duly authorized representative of Sumitomo.

WARRANTY LIMITATION:

Sumitomo warrants that this product conforms to the chemical description in the directions for use on the label, subject to the inherent risks referred to below. Sumitomo makes no other express warranties. THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS:

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY:

In no case shall Sumitomo be liable for special, indirect, or consequential damages or for any loss of profits resulting from the use of this product, and no claim of any kind shall be made against Sumitomo or its representative for the product or the extent of which the damages are not covered.

76

SUMITOMO CHEMICAL AMERICA, Inc.

345 PARK AVENUE
NEW YORK, N.Y. 10017

Net Content: 1000 g (2.2 lbs)
1000 g (2.2 lbs)

77