

5409053

 $\frac{300}{18}$

MAR 30 1992

Mr. Ronald J. White
New Mexico Department of Agriculture
Agricultural Programs and Resources
Division
Box 30005/APP
Las Cruces, NM 88003-0005

Dear Mr. White:

Subject: Sodium Fluoroacetate (Compound 1080) Livestock
Protection Collar
EPA Registration No. 39508-2
Your Submission Dated November 8, 1991

Your proposed revised technical bulletin submitted November 8, 1991, is acceptable, although we found a few minor changes that should be made. Please make these changes on your final printed labeling.

1. In last "DC" (page 3), change "hand" to "hands."
2. Delete "inert" from first sentence of first full paragraph on page 6.
3. Delete "inert" from first sentence of first full paragraph under heading "Handling collars and . . . soil" on page 9.
4. Change footnote to "Use Restriction F." to read

"2 Robert A. Forrest, Registration Division
(H7505C), Office of Pesticide Programs,
U.S. Environmental Protection Agency,
401 M St. SW, Washington, DC 20460"

We have enclosed copies of the accepted container label and technical bulletin.

BEST AVAILABLE COPY

Submit five copies of your final printed labeling before you release product for shipment under the new accepted labeling.

Sincerely yours,

Robert A. Forrest
Product Manager 14
Insecticide-Rodenticide Branch
Registration Division (H7505C)

RECEIVED

3
3

DIRECTIONS FOR USE

It is a violation of state and federal laws to use this product in a manner inconsistent with its labeling or the Compound 1080 cancellation order. Misuse may result in civil or criminal enforcement action.

DO NOT REMOVE TOXICANT FROM COLLARS. DO NOT USE TORN, DAMAGED OR LEAKING COLLARS. Dispose of damaged collars in accordance with the "Storage and Disposal" instructions on this label.

Put collars on the necks of sheep or goats in fenced pastures where coyote predation is occurring or is expected to occur. Use collars only in accordance with the User Instructions and Use Restrictions contained in the accompanying Technical Bulletin.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store Livestock Protection Collars only in original container, in a dry, locked place away from food, feed, domestic animals, and corrosive chemicals. Do not store in any structure occupied by humans.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of such materials is a violation of Federal Law.

Dispose of wastes contaminated by 1080 (carcasses, wool, hair, vegetation, soil, leather clothing, and water) under three feet of soil, at a safe location, preferably on property owned or managed by the applicator and at least one-half mile from human habitations and water supplies.

When snow or frozen ground make on-site disposal impractical, up to one cubic foot of wastes may be stored in a leakproof container, in a dry, locked place for up to 90 days.

Alternatively, contact your state pesticide Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in disposing of wastes at approved hazardous waste disposal facilities.

COLLAR DISPOSAL: Dispose of collars under three feet of soil, at a safe location, preferably on property owned or managed by the applicator and at least one-half mile from human habitations and water supplies.

CONTAINER DISPOSAL: Metal and plastic containers - Triple rinse contaminated and uncontaminated containers with water. Then puncture and dispose of contaminated containers and rinse as above.

RESTRICTED USE PESTICIDE

Collars shall be sold or transferred only by registrants or their agents, certified Livestock Protection Collar applicators. Collars may be used only by specifically certified Livestock Protection Collar applicators or by their direct supervision.

For use only in the State of New Mexico

SODIUM FLUOROACETATE (COMPOUND 1080) LIVESTOCK PROTECTION COLLAR

For use on sheep or goats to kill depredating coyotes

ACTIVE INGREDIENT - SODIUM FLUOROACETATE 1.00%
INERT INGREDIENTS* 99.00%
TOTAL 100.00%

*Contains inert Tartrazine dye as a marker

KEEP OUT OF REACH OF CHILDREN

DANGER



POISON

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting at once with an emetic such as syrup of ipecac, as directed. If emetic is not available, drink 1-2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting anything, by mouth to an unconscious person. PROMPT TREATMENT IS MEDICAL ATTENTION IMMEDIATELY.

IF ON SKIN: Wash the exposed area twice with soap and water.

IF IN EYES: Flush eyes with plenty of water for at least 15 minutes.

IF ON CLOTHING: Remove contaminated clothing and wash before reuse. Wash contaminated leather, including shoes, boots, and gloves, according to the "Pesticide Disposal" section. See disposal instruction on the label.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANUFACTURED BY: Rancher's Supply, Inc.
Box 725, Alpine, TX 79831

U.S. PAT. 3,842,806 EPA ESTABLISHMENT NO. 39508-NM
EPA REGISTRATION NO. 39508-2

NET CONTENTS: 30.4 grams (1.1 oz) per collar

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of other than indicated on the label. Buyer assumes all risks of use and of this product when such use and/or handling is contrary to label instructions.

ACCEPTED
with COMMENTS
in EPA Letter Docket
MAR 30 1992