

KEYSTONE MB-11

ACTIVE INGREDIENTS:

Disodium cyanodithioimidocarbonate 4.90%

Potassium *N*-methyldithiocarbamate 8.76%

INERT INGREDIENTS 88.34%

This product contains 1.03 lb. of active ingredients per gallon and weighs 8.88 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, give patient doses of powdered charcoal immediately or all he can swallow of raw egg white, milk, gruel, or flour and water. Then induce vomiting with salt, soap, or mustard in warm water. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

KEYSTONE MB-11 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 5.1 to 10.2 fluid ounces of KEYSTONE MB-11 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.7 to 10.2 fluid ounces of KEYSTONE MB-11 per 1000 gallons of water should be employed every 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

ACCEPTED
28 APR 1980
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRY
EPA UNDER NO. 39743-6

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in an approved landfill, or bury in a safe place. Metal containers over 30 gallons should be resealed before offering for reconditioning.

GENERAL: Consult Federal, State or Local disposal authorities for approved alternative procedures.

Manufactured by:

KEYSTONE LABORATORIES, INC.

P. O. Box 1367

Decatur, AL 35602

EPA REG. NO. 39743-6

EPA EST. NO. 39743AL01

NET CONTENTS _____

24