

K-KLEAN

ID 1 DISINFECTANT

Food Plants-Livestock Buildings-Hospitals-Veterinary Clinics
NO RINSE FOOD PLANT SANITIZER
POULTRY DRINKING WATER SANITIZER

ACTIVE INGREDIENTS:

Alpha-(P-nonylphenyl)-Omega hydroxypoly
(oxyethylene)-iodine Complex.....18.05%
(Providing 1.75% titratable iodine)

Inert Ingredients.....81.95%

KEEP OUT OF REACH OF CHILDREN

DANGER

Practical Treatment: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory, shock, respiratory depression, and convulsion may be needed.

SEE ADDITIONAL CAUTION SIDE PANEL
FOR MEDICAL, INDUSTRIAL, COMMERCIAL, AND
AGRICULTURAL USE ONLY.

Manufactured by:

K-KLEAN CHEMICAL CO. INC.
DALLAS, TEXAS 75247

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION INFORMATION

For use on hard, non-porous, inanimate surfaces (glass, stainless steel, galvanized metal, porcelain, linoleum, tile, enameled and finished wood surfaces). Use on walls, floors, appliances, furnishings, and utensils.

Prepare fresh solutions daily. Do not reuse solutions. Discard solutions that are visibly soiled. Apply the solutions by flooding, immersion, swabbing, mopping, brushing to thoroughly wet all surfaces.

FOR DISINFECTION: In Hospitals, Nursing Homes, Food Plants, Commercial Freezers, Restrooms, Restaurants, and Veterinary Clinics, Farm Premises and Poultry Houses. Use 1 oz. of this Disinfectant in 2 gallons of water for one-step cleaning and disinfecting light and moderately soiled surfaces. Heavily soiled surfaces should be cleaned and rinsed before disinfecting. For effective action, surfaces must remain wet for at least 10 minutes.

Food contact surfaces and all surfaces in edible food processing areas of Federally inspected meat and poultry plants must be rinsed with potable water before reuse when using the product at this concentration. Always carefully protect or remove food and packaging materials from the area before disinfecting.

For Farms and Poultry use, remove all animals, poultry and feed from premises. Remove all litter and manure from surfaces occupied or traversed by livestock. Empty all troughs, racks, and other feeding and watering equipment. Then, clean and disinfect as above. Ventilate buildings, coops, and other closed spaces. Do not house animals or poultry until treatment has been absorbed, set or dried.

EPA Reg. No. 39978-1

EPA Est. No. 39978-TX-01

ACCEPTED

39978-1

MAY 14 1984

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR SEVERAL PESTICIDE RESISTANCE
BY THE EPA

SANITIZATION OF MEAT AND FOOD PROCESSING EQUIPMENT AND UTENSILS: Remove gross soil flushing, scraping, or mopping. Preclean surfaces with 1 oz. of this Disinfectant per gallon of water or with an suitable cleaner. Rinse well. Sanitize with 1 oz. Disinfectant in 5 gallons of water (25 ppm titra iodine). Sanitizing solution should contact surfaces for minutes. Drain well and air dry. Rinsing is not require

HAND SANITIZATION IN MEAT AND POULTRY PROCESSING PLANTS: For sanitizing hands of handlers, rinse hands with water and immerse hands in oz. per 5 gallons of water sanitizing solution (25 titratable iodine) for 30 to 60 seconds.

DISINFECTING TOILET BOWLS AND URINALS: Re gross soils from surfaces with bowl brush. Add 1 oz. of Disinfectant to the bowl water. Use brush or swab thoroughly clean bowl surfaces, especially under the Flush commode or urinal several times and rinse bowl well.

DEODORANT: This Disinfectant destroys many odors sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab a solution of 1 ounce of disinfectant per gallon of water.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Harmful or fatal if swallowed. Corrosive. Causes damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Avoid contamination of food.

Net Contents: 1 U. S. Gallon

N CTANT

Veterinary Clinics
SANITIZER
SANITIZER

poly
.....18.05%
.....81.95%

CHILDREN
Immediately flush eyes
15 minutes. For eyes,
quantity of milk, egg
if available, drink large
damage may contrain-
dicate against circulatory,
on may be needed.
DE PANEL
COMMERCIAL, AND
ONLY.

CO. INC.
47

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION INFORMATION

For use on hard, non-porous, inanimate surfaces (glass, stainless steel, galvanized metal, porcelain, linoleum, tile, enameled and finished wood surfaces). Use on walls, floors, appliances, furnishings, and utensils.

Prepare fresh solutions daily. Do not reuse solutions. Discard solutions that are visibly soiled. Apply use solutions by flooding, immersion, swabbing, mopping or brushing to thoroughly wet all surfaces.

FOR DISINFECTION: In Hospitals, Nursing Homes, Food Plants, Commercial Freezers, Restrooms, Restaurants, and Veterinary Clinics, Farm Premises and Poultry Houses. Use 1 oz. of this Disinfectant in 2 gallons of water for one-step cleaning and disinfecting light and moderately soiled surfaces. Heavily soiled surfaces should be cleaned and rinsed before disinfecting. For effective action, surfaces must remain wet for at least 10 minutes.

Food contact surfaces and all surfaces in edible food processing areas of Federally inspected meat and poultry plants must be rinsed with potable water before reuse when using the product at this concentration. Always carefully protect or remove food and packaging materials from the area before disinfecting.

For Farms and Poultry use, remove all animals, poultry and feed from premises. Remove all litter and manure from surfaces occupied or traversed by livestock. Empty all troughs, racks, and other feeding and watering equipment. Then, clean and disinfect as above. Ventilate buildings, coops, and other closed spaces. Do not house animals or poultry until treatment has been absorbed, set or dried.

EPA Reg. No. 39978-1

EPA Est. No. 39978-TX-01

ACCEPTED
39978-1
MAY 14 1984
UNDER THE FEDERAL INSECTICIDE
ACT AND THE FEDERAL FOOD, DRUG AND
COSMETIC ACT FOR SUBSTANCES REGISTERED
BY EPA REG. NO.

SANITIZATION OF MEAT AND FOOD PROCESSING EQUIPMENT AND UTENSILS: Remove gross soils by flushing, scraping, or mopping. Preclean surfaces with 1 oz. of this Disinfectant per gallon of water or with another suitable cleaner. Rinse well. Sanitize with 1 oz. of Disinfectant in 5 gallons of water (25 ppm titratable iodine). Sanitizing solution should contact surfaces for 1-2 minutes. Drain well and air dry. Rinsing is not required.

HAND SANITIZATION IN MEAT AND POULTRY PROCESSING PLANTS: For sanitizing hands of food handlers, rinse hands with water and immerse hands in a 1 oz. per 5 gallons of water sanitizing solution (25 ppm titratable iodine) for 30 to 60 seconds.

DISINFECTING TOILET BOWLS AND URINALS: Remove gross soils from surfaces with bowl brush. Add 1 oz. of this Disinfectant to the bowl water. Use brush or swab to thoroughly clean bowl surfaces, especially under the rim. Flush commode or urinal several times and rinse bowl brush well.

DEODORANT: This Disinfectant destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails, refuse containers, etc.), swab on a solution of 1 ounce of disinfectant per gallon of water.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Harmful or fatal if swallowed. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Avoid contamination of food.

Net Contents: 1 U. S. Gallon

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	MAY 14 1984
	NAME OF PESTICIDE PRODUCT	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>		
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		
[] [] [] []		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE