



GOSSETT

R. O. L. ~~XXXXXXXXXX~~ PREMIX

(Contains RABON[®] Oral Larvicide)

To prevent the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	7.76%
INERT INGREDIENTS	92.24%*
TOTAL	100.00%

*Refers only to ingredients which are not larvicidal.

40034-1

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Net Wt. 45 lbs.

EPA Reg. No. 40034

EPA Estab. No. 40034-OH-01

RABON is the registered trademark of Shell Oil Company. U.S. Patent No. 3,102,842

Manufactured by

~~XXXXXXXXXX~~
GOSSETT NUTRITION, INC.
P. O. Box 512
MARION, OHIO 43302

GOSSETT R.O.L. PREMIX

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When fed to cattle, RABON® Oral Larvicide passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, RABON® Oral Larvicide helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

RABON® Oral Larvicide Premix will mix uniformly in cattle feeds when standard mixing procedures are followed. Thus usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution of RABON® 7.76 Oral Larvicide Premix throughout the feed mix. The premix should be preblended with ground grain, protein supplements, mineral mixes, etc., before being added to roughages, such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if RABON® 7.76 Oral Larvicide Premix is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixers, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with RABON® 7.76 Oral Larvicide Premix should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, RABON® 7.76 Oral Larvicide Premix should not be mixed in liquid feed supplements.

In a Concentrate Feed—Roughage Fed Separately

RABON® 7.76 Oral Larvicide Premix can be used to prepare concentrate feeds that will provide 792 mg. of RABON per animal per day.

To prepare a larvicidal concentrate feed, mix RABON® 7.76 Oral Larvicide Premix according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

MIXING DIRECTIONS

Pounds of Concentrate Consumed per Animal per Day	RABON in the Concentrate mg/lb	Pounds of RABON® 7.76 Oral Larvicide Premix per Ton of Concentrate
5	1584 0.35	900
10	792 0.18	450
15	528 0.12	300
20	396 0.087	225
50	159 0.035	90
100	79 0.018	45
150	53 0.012	30
200	39 0.0087	23
250	32 0.0069	18
300	28 0.0059	15

Feed the appropriate larvicidal concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200-1700 pounds, increase the amount of premix per ton of concentrate to 1 1/2 times that recommended.

Start feeding RABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Rations containing RABON® Oral Larvicide may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment.

RABON® Oral Larvicide prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle but is not effective against existing adult flies.

In some cases supplemental fly control measures may be needed in and around cattle lots and barns to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, RABON® Oral Larvicide should be used in conjunction with other good management and sanitation practices.

In a Complete Ration—No Other Roughage Fed

RABON® 7.76 Oral Larvicide Premix can be used to prepare rations that contain 26.4 mg. of RABON per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of RABON® 7.76 Oral Larvicide Premix per ton of complete mixed ration containing both grain and roughage.

Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds or to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticides that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

<p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (MH-567) WASHINGTON, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i></p>	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP code)</p> <div style="text-align: right; padding-right: 50px;"> <p>161/85431 181 1</p> </div>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p><input type="checkbox"/> ATTACHMENT IS APPLICABLE</p>		
SIGNATURE OF APPROVING OFFICIAL		DATE