

ROSSIN RATION NO. 10

NEW BARI-FLYER
50% RATION (CATTLE LACTATING)

To prevent the development of Horn Flies, Face Flies, House Flies, and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphite	1.00%
INERT INGREDIENTS	99.00% *
TOTAL	100.00%

* Refers only to ingredients which are not larvicidal

GENERAL ANALYSIS

Calcium (Ca), Maximum	13.00%
Calcium (Ca), Minimum	11.00%
Phosphorus (P), Minimum	6.00%
Salt (NaCl), Maximum	26.50%
Salt (NaCl), Minimum	23.50%
Iodine (I), Minimum	0.007%
Magnesium (Mg), Minimum	8.00%
Iron (Fe), Minimum	0.20%
Copper (Cu), Minimum	0.02%
Cobalt (Co), Minimum	0.01%
Manganese (Mn), Minimum	0.16%
Zinc (Zn), Minimum	0.20%
Selenium (Se)	0.0008%
Vitamin A, Minimum USP Units per lb.	350,000
Vitamin D-3, Minimum USP Units per lb.	200,000
Vitamin E, Minimum USP Units per lb.	50

ACCEPTED
With comments of
EPA letter dated
AUG 31 1982
EPA Reg. No. 40037-3

BEST DOCUMENT AVAILABLE

INGREDIENTS

Monocalcium phosphate, dicalcium phosphate, calcium carbonate, salt, yeast culture, corn distillers dried grains with solubles, dried molasses, vitamin A acetate, D-activated animal sterol (source of vitamin D-3), vitamin E supplement, magnesium oxide, copper oxide, cobalt carbonate, ethylenediamine dihydrochloride (EDDI), iron sulfate, manganous oxide, zinc oxide, iron oxide, mineral oil, artificial flavoring, sodium selenite, potassium sulfate, and magnesium sulfate.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See reverse of tag for additional precautionary statements.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION
Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

Net Wt. 50 lbs.

EPA Reg. No. 40037
EPA Estab. No. 10037-01-01

RABOL is the registered trademark of Shell Oil Company.

U.S. Patent No. 2,102,842

Manufactured By
ROSSIN RATION, INC.
P.O. Box 512
MADISON, OHIO 43002

REGISTRATION NO. 1478
**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Allow free choice to cattle. Cattle should consume an average of $\frac{1}{2}$ lb. of the mineral per 100 pounds of body weight per month. This will supply the recommended average daily intake of 70 mg. of organo-arsenic per hundred pounds body weight. Animals should not be starved for minerals before initiation of the RABON Cattle Mineral Mix program but stop all salt from other feed after the program is started. Do not feed other free choice sources of salt or mineral containing salt in any form. Allow one mineral feeder for each 10-20 head, and place near watering and loafing areas. Put out no more than one month's supply at a time, and protect from rain.

If consumption of Beef Pre-Mix No. 14 is above $\frac{1}{2}$ lb. per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below $\frac{1}{2}$ lb. per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Beef Pre-Mix No. 14 early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Beef Pre-Mix No. 14 may be fed up to 30 days after and to lactating dairy cows.

Beef Pre-Mix No. 14 prevents the development of horn flies, face flies, house flies, and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.


Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Beef Pre-Mix No. 14 should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

BEST DOCUMENT AVAILABLE



subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)