

ACCEPTED

40080-2
MAR 27 1981

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
25 U.S.C. 135
Registration No. 10000
EPA Reg. No. 40080-2

SODIUM HYPOCHLORITE SOLUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive, may cause severe skin irritation or chemical burns to broken skin. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:
STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with gross filth such as feces, urine, etc., or with ammonia, acids, detergents or other chemicals may release hazardous gases irritating to eyes, lungs and mucous membranes.

manufactured by

**OUTDOOR WORLD
DISTRIBUTORS, INC.
6900 N.W. 12th Avenue
Ft. Lauderdale, FL 33309**

EPA REG. NO. 40080-2 EPA EST. NO.
NET CONTENTS

DIRECTIONS GENERAL CL

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THE LABEL DIRECTIONS.

DIRECTIONS FOR SWIMMING POOL CHLORINATION AND SANITIZING:
For a new pool or for spring start-up, add this sodium hypochlorite solution for each gallon of water to 5 ppm available chlorine by weight.
For Pool Maintenance:
A) Adjust pool water pH to 7.6 - 7.8 range.
B) Add manually or by a feeder device to maintain an available chlorine residual of 1.0 ppm available chlorine by weight. 1.0 ppm available chlorine will depend on pool size. Use test kit to make certain the pH is 7.6 - 7.8.

DIRECTIONS FOR DISINFECTING FOR HOME WELL:
Dilute this sodium hypochlorite solution to 10 parts softened water. Mix thoroughly and begin feeding of solution with a free available chlorine residual of 5.0 throughout the distribution system, as determined by bacteriological sampling must be conducted at the National Interim Primary Drinking Water Act with a DPD chlorine test kit.
*Contact your local Health Department for more information.

DIRECTIONS FOR SANITIZING FOOD:
Clean equipment in the normal manner. Apply this sodium hypochlorite solution to obtain contact with disinfectant for a minimum of 10 minutes. Rinse with water. After treatment with sodium hypochlorite, one fluid oz. (2 tbsps) of this sodium hypochlorite solution will provide approximately 200 ppm available chlorine.

NOTE: This product degrades with age. As necessary to obtain the required amount of available chlorine, use a test kit.
STORAGE AND DISPOSAL: Store in original container, away from direct sunlight. In case of spill, flood empty container thoroughly with water and place in trash collection or burying. Do not use if container is damaged. Do not contaminate food or feed by storage.

"FOR SWIMMING POOL CHLORINATION AND SANITIZING."

ACTIVE INGREDIENT: SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

ACCEPTED

40080-2
MAR 27 1981

U.S. Dept. of Agriculture
Federal Insecticide, Fungicide, and Rodenticide Act

SODIUM HYPOCHLORITE SOLUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive, may cause severe skin irritation, chemical burns to broken skin. Causes eye damage. Do not get on eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash thoroughly. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until odors have dissipated.

ENVIRONMENTAL HAZARDS: This pesticide is highly toxic. Keep out of lakes, streams or ponds. Treated effluents should not be discharged into lakes, streams, ponds or public waterways without a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS:

STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with organic materials such as feces, urine, etc., or with ammonia, acids, or other chemicals may release hazardous gases. Avoid contact with eyes, lungs and mucous membranes.

manufactured by
OUTDOOR WORLD DISTRIBUTORS, INC.
6900 N.W. 12th Avenue
Ft. Lauderdale, FL 33309

EPA REG. NO. 40080-2 EPA EST. NO. 1000000000
NET CONTENTS

"FOR SWIMMING POOL CHLORINATION AND SANITIZING."

ACTIVE INGREDIENT
SODIUM HYPOCHLORITE 9.2%
INERT INGREDIENTS 90.8%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID: If on skin, wash with plenty of soap and water. If in eyes, flush with water for at least 15 minutes. Get medical attention. If swallowed, drink large quantities of milk, or gelatin solution or, if these are not available, drink large quantities of water. Do NOT give vinegar or other acids. Do NOT induce vomiting. Get prompt medical attention.

See additional precautions on side panel

DIRECTIONS FOR USE GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

DIRECTIONS FOR SWIMMING POOL CHLORINATION

For a new pool or for spring start-up, superchlorinate with one (1) pint (16 fluid ozs) of this sodium hypochlorite solution for each 2500 gallons of water. This dosage is equivalent to 5 ppm available chlorine by weight.
For Pool Maintenance:
A) Adjust pool water pH to 7.6 - 7.2 range and maintain.
B) Add manually or by a feeder device this sodium hypochlorite solution at a rate to maintain an available chlorine residual of 0.8 to 1.0 ppm. One-half (1/2) pint (8 fluid ozs) of this sodium hypochlorite solution for each 6000 gallons of water will provide 1.0 ppm available chlorine by weight. Frequency of additions to maintain 0.8 to 1.0 ppm available chlorine will depend upon temperature and number of swimmers. Use test kit to make certain the pH and chlorine residual are in the proper range.

DIRECTIONS FOR DISINFECTION OF POTABLE WATER FOR HOME WELL WATER SYSTEMS

Dilute this sodium hypochlorite solution in the ratio of one part of sodium hypochlorite solution to 9 parts softened water. Mix sodium hypochlorite solution and water thoroughly and begin feeding of solution with a hypochlorinator (metering pump). Maintain a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm throughout the distribution system, as determined by a DPD chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Check water frequently with a DPD chlorine test kit. Contact your local Health Department for further details.

DIRECTIONS FOR SANITIZING FOOD PROCESSING EQUIPMENT

Clean equipment in the normal manner. Just before using, rinse all surfaces thoroughly with this sodium hypochlorite solution containing 200 ppm available chlorine. Maintain contact with disinfectant for a minimum of two minutes. Do not rinse with water after treatment with sodium hypochlorite solution. Do not eat overnight. Addition of one fluid oz (2 tblsp) of this sodium hypochlorite solution per 2 gallons of water will provide approximately 200 ppm available chlorine by weight.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage as necessary to obtain the required level of available chlorine.
STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight. In case of spill, flood area with large quantities of water. Rinse empty container thoroughly with water and either return to manufacturer or discard by placing in trash collection or burying in an approved landfill. Product or rinsate that cannot be used should be diluted with water and disposed of in a sanitary sewer. Do not contaminate food or feed by storage, disposal or cleaning of equipment.