

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations

For sanitizing food processing equipment, dishes, equipment, and utensils, silverware, glasses, sink tops, countertops, refrigerated storage equipment and other hard surfaces. **No potable water rinse is required.**

Wash and rinse all articles thoroughly, then apply a solution of O-C SAN-Q No. 1 in 4 gallons of water. (150ppm active)

Surfaces should remain wet for at least one minute following application and drying. Fresh solution should be prepared daily or when the solution becomes visibly dirty. For mechanical application, use solution spray not be used for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. **No potable water rinse is required.**

Dishes, silverware, glasses, cooking utensils and other similar food processing equipment can be sanitized by immersion in a 1 oz. 4 gallon solution of O-C SAN-Q No. 1. **No potable water rinse is required.**

At 1 oz./4 gallons, O-C SAN-Q No. 1 fulfills the criteria of Appendix E of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendation of the U.S. Public Health Services in waters up to 1000ppm of hardness, calculated as calcium when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. O-C SAN-Q No. 1 in 4 gallons of warm water. **No potable water rinse is required.**

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements

Hazards to humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED WITH STORAGE

OR DISPOSAL.

OPEN DUMPING IS PROHIBITED.

DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL

PESTICIDE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE OR EQUIVALENT AND DEPOSE IN A LANDFILL OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE, OR LOCAL DISPOSAL REGULATIONS FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LINDEN OPEN BURNING.



O-C SAN-Q No. 1

Disinfectant Sanitizer Fungicide
Deodorizer for Hospital, Institutional,
Industrial, School, Dairy and Other
Farm and Home Use

ACTIVE INGREDIENTS

Didecyl dimethyl ammonium chloride 7.5%

Isopropanol 3.0%

INERT INGREDIENTS 89.5%

100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

See Left Panel for Additional Precautionary Statements.

EPA Registration No. 10157-1

Net Contents

gal. BATCH NO.



O.C. SAN-Q No. 1

Disinfectant-Sanitizer-Fungicide
Deodorizer for Hospital, Institutional,
Industrial, School, Dairy and Other
Farm and Home Use

INGREDIENTS	
Dimethyl ammonium chloride	7.5%
Alcohol	3.0%
INGREDIENTS	89.5%
	100.0

**KEEP OUT OF REACH OF CHILDREN
DANGER**

Statement of Practical Treatment

Contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before use.

If drink promptly a large quantity of milk, egg whites, gelatin solution, or fruit juice. If not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

PHYSICIAN: Probable mucosal damage may contraindicate the use of antacids. Measures against circulatory shock, respiratory depression, and other complications may be needed.

See label for Additional Precautionary Statements

Registration No. 10457-1
Net Weight 1 gal. BATCH NO.

Directions for Use GENERAL CLASSIFICATION

For disinfection of floors, walls, ceilings, and other hard surfaces in hospital, institutional, industrial, school, dairy and other farm and home use. For disinfection of poultry equipment, animal quarters and kennels.

ADD 2 oz. of O.C. SAN-Q No. 1 to 4 gallons of water. Apply to hard surfaces with mop or cloth.

At 2 oz. / 4 gallon use-level, O.C. SAN-Q No. 1 is effective against Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton mentagrophytes (the athlete's foot fungus).

Disinfection of Institutions, Industries, Schools and Homes

For disinfection of floors, walls, ceilings, and other hard surfaces in hospital, institutional, industrial, school, dairy and other farm and home use.

Add 2 oz. of O.C. SAN-Q No. 1 to 4 gallons of water. Apply to hard surfaces with mop or cloth.

At 2 oz. / 4 gallon use-level, O.C. SAN-Q No. 1 is effective against Staphylococcus aureus, Salmonella choleraesuis, and Trichophyton mentagrophytes (the athlete's foot fungus).

Disinfection of Barber Tools

Pre-cleaned barbers' tools (clippers, combs, brushes, razors, and scissors) can be disinfected by immersion in a 2 oz. / 4 gallon solution of O.C. SAN-Q No. 1.

Disinfection of Poultry Equipment, Animal Quarters and Kennels

Poultry brooders, watering tanks, feeding equipment and other animal quarters (such as pens and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. O.C. SAN-Q No. 1 to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Manufactured By:

OMNI-CHEM COMPANY, INC.

1889 MT. DIABLO STREET
CONCORD, CALIFORNIA 94522

FOR REORDER CALL: (415) 825-0300