

13 / # 7-1-7

GENERAL DIRECTIONS

Use the following recommended dosages per acre in sufficient water for good coverage, usually 100 gals. For Aircraft Application: Minimum quantity of water: Approximately 100 gals. per acre, consistent with crop growth and good coverage.

NOTE: Do not apply to Alfalfa or Clover within 7 days of harvest or grazing. Do not apply during full bloom.

ALFALFA RECOMMENDED APPLICATION

ALFALFA WEEVIL LARVAE Hay Crop (West): Treat in May or June using 2.0 qts. per acre. Hay Crop (East): Treat in Spring when most of the tips are beginning to show feeding, using 2.0 qts. per acre. A second application may be necessary. If harvest infestation is severe, treat stubble as soon as crops are removed.

ALFALFA CATERPILLAR Use 1 1/2 qts. per acre. Treat when alfalfa starts to show signs of injury. Caterpillars per sweep net per acre usually when alfalfa is approximately 1.5 qts.

CLOVER LEAF WEEVIL Use 1 1/2 qts. per acre. Treat in Spring when alfalfa is 2 to 3 inches tall.

GRASSHOPPERS Use 1.1 qts. per acre. Treat after cutting and before alfalfa plants are 10 inches tall. A second application may be necessary.

MEADOW SPITTLERBUG Spring Treatment: Use 1 qt. per acre. Apply within a week after eggs begin to hatch. Fall Treatment: Use 1 qt. per acre. Apply in early September.

PEA APHID on HAY or SEED CROP Use 2 qts. per acre. Treat before bloom or in even bloom stage.

POTATO LEAFHOPPERS Use 1 1/2 qts. per acre. Apply when alfalfa is about 1/3 grown or earlier if insects become abundant.

SPOTTED ALFALFA APHID Use 1 1/2 qts. per acre. Treat stem-ting when 1 qt. per plant of alfalfa seedlings stand when alfalfa is per acre stage.

ACCEPTED
AUG 06 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 4715-94

70831-4
**ALFALFA and CLOVER
INSECT CONTROL**

Chlorpyrifos, Methoxyfenosulfuron and Deltamethrin per gallon.

ACTIVE INGREDIENTS	By Weight
Methoxyfenosulfuron	24.0
Chlorpyrifos	24.0
Pyrethrin	4.0
Asymetric Petroleum Derivative Solvent	47.0
INERT INGREDIENTS	5.0
	Total: 100.0

*Equivalent to 21.2 gals. of methoxyfenosulfuron, 1.1 gal. of chlorpyrifos and 1.0 gal. of other solvent and inert ingredients.

CAUTION Keep out of Reach of Children

See side panel for complete caution statement.

NET CONTENTS: 1.0 GAL. (128 FL. OZ.)
COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

CLOVER RECOMMENDED APPLICATIONS

CLOVER LEAF WEEVIL Use 1 1/2 qts. per acre. Treat in Spring when clover is 2 to 3 inches tall.

GRASSHOPPERS Use 1 1/2 qts. per acre. Treat when grasshoppers are young. A second application may be necessary.

MEADOW SPITTLERBUG Spring Treatment: Use 1.2 qts. per acre, treat within a week after eggs begin to hatch. Fall Treatment: Use 1.2 qts. per acre, treat in early September.

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing fumes or spray mist. Avoid getting in eyes on skin or on clothing. In case of contact, wash with soap and water. For eyes, get medical attention. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy containers by breaking or crushing and bury in non-cropland away from water supplies.

NOTICE: Seller warrants this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 4715-94 EPA Est. 4715-CO-1

BEST AVAILABLE COPY

COMPANY
NUMBER 788-

PRODUCT
SERIAL NO. 18

LABEL
APPROVAL
DATE 5-10-85

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	781-11	5/3/85
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended.</i>	NAME OF PESTICIDE PRODUCT	
	Fungicide, Transparent Emulsion Spray	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[Faded Name]
 [Faded Name]
 [Faded Name]

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

1. Submit to the Administrator, at the address shown on the label, all data required for registration or reregistration of your product under 40 CFR 152.23(a)(1) when the following requirements are met:

2. Submit to the Administrator, at the address shown on the label, all data required for registration or reregistration of your product under 40 CFR 152.23(a)(1) when the following requirements are met:

3. Label the label with the following information before you release the product for shipment:

a. Add the phrase "EPA Registration No. 781-11."

b. Revise your Environmental Hazards Paragraph to read:

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water. Run-off from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate soils. Do not contribute water by cleaning of equipment or disposal of wastes.

c. Revise your pesticide disposal directions to read:

Wastes resulting from the use of this pesticide may be disposed of or stored at an approved water treatment facility.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
[Signature]	5/3/85

Revise your Container Disposal directions to read:

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

- d. Revise the If Swallowed section of your Statement of Practical Treatment to read:

Drink 1 or 2 glasses of water and induce vomiting by sticking finger down throat.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

JOB:92902:Ellenberger:RD85-15:5/2/85:Del:5/10/85:Kendrick:898-1270:DAA

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash hands after using.

In food handling areas, application is limited to crack and crevice and/or spot treatment. Do not apply directly to food or feedstuffs. In commercial food handling facilities, cover or remove any food or food processing equipment during application. Do not apply while food processing is underway. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Get immediate medical attention. If in eyes: Flush with plenty of water for at least 15 minutes. Get medical attention. If on skin: Flush with plenty of water for at least 15 minutes.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition appear, atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL/CHEMICAL HAZARDS

Do not apply this water-based product to conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store this water-based formulation at temperatures below 32°F (0°C).

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures.

CONTAINER DISPOSAL: For Containers One Gallon And Under: Do not use empty container. Wrap container and place in trash collection. For Containers Over One Gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and Local procedures.

FLEATHAL

TRANSPARENT EMULSION** SPRAY

A COMBINATION OF INSECTICIDES OFFERING RESIDUAL KILL OF COCKROACHES FOR UP TO 4 WEEKS

For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), Clover mites, centipedes, millipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, house flies, mosquitoes, gnats, flying moths, confused flour beetles, sawtoothed grain beetles, shiny spider beetles, grain mites, red flour beetles, Angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, clothes moths.

ACTIVE INGREDIENTS:

- *† Resmethrin[5-(phenylmethyl)-3-furanyl]methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate 0.05%w/w
- * Chlorpyrifos[0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)phosphorothioate].....

INERT INGREDIENTS:.....

* Cis:trans isomers ratio: max. 30%(cis) and min. 70%(trans)
 † U.S. Patent Nos. 3,465,007 and 3,542,928
 * Perick's SBP-1382® brand of resmethrin insecticide.
 Perick Corp., 1050 Wall Street West, Lyndhurst, N.J. 07071
 * U.S. Patent No. 3,244,586 Dursban is a Trademark of Dow Chemical Co.
 ** U.S. Patent No. 3,683,078

CONTAINS DURSBAN * INSECTICIDE

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que le explique la etiqueta la haya sido explicada ampliamente. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

In food handling establishments, can be used in non-food areas as a contact or residual spray. In FOOD AREAS, application is limited to spot and/or crack and crevice treatment.

E.P.A. REG. No.

E.P.A. EST. No. 788 PA-1

NET CONTENTS:

Distributed by: Victory Chemical Co. Philadelphia, PA

**TESTED BY
VICTORY CHEMICAL CO.
IN 1978 DURS-45%**

100.00% w/w

MAY 10 1985

488/18

DEPT. OF HEALTH 6011

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE IN NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF HOTELS, MOTELS, HOMES, PRESCHOOLS, APARTMENT BUILDINGS, COMMERCIAL, OFFICE, INSTITUTIONAL AND STORAGE BUILDINGS, RECREATIONAL BUILDINGS, BOATS, SHIP CABINS, AND HOSPITALS. IN HOSPITALS, REMOVE PATIENTS PRIOR TO TREATMENT. VENTILATE ROOMS FOR TWO HOURS BEFORE RETURNING PATIENTS.

THIS PRODUCT CAN BE USED IN NON FOOD AREAS SUCH AS CORRIDORS, OFFICES, FOYERS, LAVATORIES, GARAGES, UTILITY ROOMS, STORAGE ROOMS AND BASEMENTS, OF PACKAGING AND PROCESSING PLANTS, MILK HANDLING AREAS OF MILK PLANTS, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOUR AND FEED MILLS, AND BAKERIES.

IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, APPLICATION IS LIMITED TO SPOT AND/OR CRACK AND CREVICE TREATMENT.

ALSO, FOR USE IN AND AROUND CAMPERS, MOBILE HOMES, BUSES AND AUTOMOBILES. THIS PRODUCT CAN BE USED IN CARGO PLANES, TRUCKS, BOXCARS AND SHIP CARGO HOLDS ONLY IF THE VEHICLES ARE NOT USED FOR TRANSPORT OF FOOD OR FOOD PRODUCTS AT ANY TIME. ALSO, FOR USE IN DOG KENNELS AND HORSE STABLES. DO NOT SPRAY ON HORSES AND DOGS. DO NOT SPRAY DOG AND HORSE FOOD OR FEED ITEMS.

NOTE TO PEST CONTROL OPERATIONS: One gallon (3.79 L) of spray is sufficient to treat 1500 linear feet (457 m) with a 6 inch (15.2 cm) spray band.

For contact and residual sprays, use this product without diluting. For repeat applications, dilute with equal parts of water.

CONTACT SPRAY-KILLS AND REDUCES INFESTATIONS:

Cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, centipedes, millipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, confused flour beetles, sawtoothed grain beetles, shiny spider beetles, grain mites, red flour beetles, Angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Use as a spot treatment. When possible, spray directly on

pests to reduce infestations. Apply as a coarse spray or with a paint brush into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators and other appliances, baseboards and storage areas. Apply spray until areas are wet, but not to the point of run-off. Repeat as necessary.

ANTS (Including Fire Ants): To kill ants, spray trails, nests and points of entry within buildings and homes. Spray directly on ants where possible. Repeat as necessary.

FLEAS, BROWN DOG AND LONE STAR TICKS: Thoroughly apply as a spot treatment to infested areas such as pets' beds and resting quarters; nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean fresh bedding after treatment of pet areas. **DO NOT TREAT PETS WITH THIS PRODUCT.** Treat pets with a product registered for animal application such as SBP-1382/Bioallethrin Aqueous Pressurized Spray, EPA Reg. No. 432-482, to control the source of flea and tick infestation.

CARPET BEETLES AND CLOTHES MOTHS: Do not spray clothing with this product. Do not allow clothing to contact areas where this product has been used. Spray edges of carpeting and under carpeting rugs, and closets. Make localized applications to the floor and baseboards. Spray directly into cracks, crevices, closets, and infested areas of shelving and/or wherever these insects are seen or suspected. Repeat as necessary.

RESIDUAL SPRAY: AIDS IN THE CONTROL OF COCKROACHES, ROACHES (Waterbugs), CRICKETS, BOXELDER BUGS, SCORPION BEETLES, SILVERFISH (Firebrats), CENTIPEDES, MILLIPEDES, EARWIGS, SOWBUGS (Pillbugs). Kills up to 4 weeks. Apply on baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in kitchens and bathrooms. In basements, apply in hiding places, in cracks and crevices, lower walls and dark places. Spray surfaces until wet, but not to the point of run-off.

PESTS OUT-OF-DOORS

DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (Lawns, Flowering Plants, Shrubs, Trees, Evergreens, etc.) BECAUSE OF POSSIBLE INJURY.

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use this product to control ants, clover mites, crickets, earwigs, millipedes

and sowbugs (pillbugs) on the outside surfaces. help prevent infestation of buildings treat the burrow to a height of 2 to 3 feet (0.5 to 1 m), where pests are found entrance. Also, apply as a residual spray to outside buildings, including porches, screens, window panes, garages, refuse dumps and other areas where they congregate or have been seen. Repeat treatment to maintain effectiveness.

HOUSE FLIES, MOSQUITOES, GNATS, FLEAS, OUTDOORS: For outdoor use only as an annoyance from these insects. Spray outside surrounding doors, window frames or wherever these insects congregate. Repeat as necessary.

ANTS, BROWN DOG TICKS, CRICKETS, FLEAS, OUTDOORS: For treatment of localized infestations in outdoor areas where there are weeds or other areas, spray infested areas thoroughly. For ants, spray hills and runways. Repeat application as infestation as reinfestations occur.

TO CONTROL FIRE ANTS: Dilute 1.5 pints (0.71 L) in 1 gallon (3.79 L) of water. Apply 1 gallon (3.79 L) gentle rain to each fire ant mound using a watering can. Thoroughly wet the mound and surrounding area to a diameter (1.22 m). For best results, apply in cool weather (18.5°C-27°C), or in early morning or late evening. Do not disturb the mounds as they appear. Do not disturb the treatment. If possible, treat all mounds in the vicinity as reinfestations occur.

Larger quantities of solution can be prepared for large scale treatments. Equipment to deliver the diluted product should be used at the rate of 1 gallon (3.79 L) per mound.

WARRANTY STATEMENT

Our recommendations for use of this product are believed to be reliable. The use of this product for the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained. The buyer must assume all responsibility, including damage, resulting from its misuse as such, or in conjunction with other materials.