

PM
408

FOR USE ON	TO CONTROL	In 100 Gal.	In 1 Gal.	COMMENTS
Apples, Pears	European Red Mite	1-2 pts.	2 tsp.	Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest.
	Clover Mite			
	Two Spotted Mite			
	Woolly Apple Aphid	1 pt.	1 tsp.	
	Forbes Scale			
	Green Apple Aphid	1 1/2 pts.	1 1/2 tsp.	
	Rosy Apple Aphid			
Do not exceed 15 lbs. actual Malathion per acre.				
Tomatoes	Spider Mites	1 1/2 pts.	1 1/2 tsp.	Make one or more applications as needed. Do not use within 3 days of harvest.
	Aphids			
	Do not exceed 2 1/2 lbs. actual Malathion per acre.			
Beans	Mexican Bean Beetle	2 pts.	2 tsp.	
	Spider Mites, Leafhopper			
Do not exceed 1 1/2 lbs. actual Malathion per acre				
Peas	Pea Aphid	2 pts.	2 tsp.	Make one or more applications. Do not use within 3 days of harvest. Do not use pea vines for forage.
Do not exceed 2 1/2 lbs. actual Malathion per acre.				
Potatoes	Aphids	2 pts.	2 tsp.	
Do not exceed 3 lbs. actual Malathion per acre.				
Cabbage	Aphids	2 pts.	2 tsp.	Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on Broccoli and Turnips, 7 days on others.
Broccoli	Imported Cabbage			
Kale	Worm			
Mustard				
Turnip				
Do not exceed 2 1/2 lbs. actual Malathion per acre.				
Flowers	Oyster Shell Scale	1 pt.	1 tsp.	Apply in summer when infestation occurs. Repeat in 10-12 days. For scales, apply when crawlers have settled on foliage.
Ornamentals	Eunonymus Scale			
Shrubs	Aphids, Spider Mites	1 1/2 pts.	1 1/2 tsp.	
	White Fly, Thrips, Mealy Bugs, Japanese Beetle Adult, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Bagworms, Juniper scale	2 pts.	2 tsp.	
	Soft Brown Scale	2 1/2 pts.	2 1/2 tsp.	
	Pine Needle Scale	4 pts.	4 tsp.	
CAUTION: Do not use on certain ferns, including Boston Maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.				
GENERAL DIRECTIONS FOR ALFALFA AND CLOVER				
To control various insects on Alfalfa and Clover use the following recommended dosage per acre in 3-10 gallons of water for good coverage. Do not apply during full bloom. Malathion may be applied on Alfalfa and Clover with no time limitations providing that no more than 1 1/2 lbs. actual Malathion per acre is used. Note: One quart of 57% Malathion contains 1 1/4 lbs. actual Malathion.				

ACCEPTED

MAY 15 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 40831-34

FALLS C

57% MA

EMULSIFIAB

(Made With P

For Control of

W

KEEP OUT O

See Side Pa

ACTIVE INGREDIENTS:

Malathion*

Xylene

INERT INGREDIENTS:

* O, O-dimethyl dithiophosphate of di...

gallon.

CONTENT

EPA Reg. No. 40831-34

FALLS CHEMICALS, INC.



CHEMICALS, INC.

MALATHION

ABLE CONCENTRATE

(with Premium Grade Malathion)

of Animal and Plant Pests

WARNING:

OF REACH OF CHILDREN

Panels for Additional Warnings

By Weight

.....	57%
.....	34%
.....	9%
Total	100%

diethyl mercaptosuccinate. Contains 5 pounds Malathion per

ENTS: ☐ ☐ ☐ ☐ GALLONS

EPA Est. No. 40831-MT-03

RECOMMENDED APPLICATIONS FOR ALFALFA

ALFALFA WEEVIL LARVAE: On Hay Crop in West—Treat alfalfa in May or June using 1 quart per acre.

On Hay Crop in East—Treat in Spring when most of the tips are beginning to show using 1½ pints per acre. A second application may be necessary. If harvest is severe treat stubble as soon as crop is removed.

CLOVER LEAF WEEVIL: Treat in Spring when alfalfa is 2-6 inches high using 1 pint per acre.

GRASSHOPPERS: Treat after cutting and before plants are 6 inches tall using 1 pint per acre. A second application may be necessary.

PEA APHID ON HAY OR SEED CROP: Treat before bloom or in evening during bloom using 1 - 1½ pints per acre.

POTATO LEAFHOPPERS: Treat when alfalfa is about half grown or earlier if insects are abundant, using 1 pint per acre.

SPOTTED ALFALFA APHID: Treat seedlings when ½-1 aphid per plant is found or stands when 5 aphids per leaf are found. Use 1 pint per acre.

RECOMMENDED APPLICATIONS FOR CLOVER

CLOVER LEAF WEEVIL: Treat in Spring when clover is 2-6 inches tall, using 1 pint per acre.

GRASSHOPPERS: Treat when grasshoppers are young. A second application may be necessary. Use 1½ pints per acre.

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry Houses, Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn meal and add to 2 gallons of water. Paint or spray on desired surfaces.

POULTRY: To control Chicken Lice add 1 oz. to 1 gallon water and apply directly to birds. This 1 gallon of diluted mixture should be applied to 100 to 500 birds. To spray walls and litter make a solution of 2 oz. in 1 gallon water. Do not apply this higher dosage. As a roost paint for lice control mix 2 ounces of 57% Malathion in one quart of water and apply at the rate of one pint of finished spray per 150 feet of roost.

BEEF CATTLE: For control of Lice, add 1 gallon to 100 gallons of water. Apply as a coverage spray. Repeat applications at two week intervals if needed. For control of ticks apply 1 to 2 gallons per 100 gallons of water. Apply a complete coverage spray. Repeat applications at two week intervals if needed.

SWINE: For control of Lice, apply 1 gallon per 1000 gallons of water. Apply a complete coverage to animals, pens and litter. One treatment may be sufficient. Repeat applications if needed.

BACK RUBBING DEVICES: For the reduction of Lice on Beef Cattle, apply a 2% solution (1 pint in 4.3 gals. of fuel oil or kerosene) to back rubbing device. There may also be a problem in Horn Flies. Make rubbing device accessible one to 35 to 45 head of cattle. Re-treat every 2 or 3 weeks. **DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS**

CAUTION: Do not apply to dairy animals. Do not treat calves under one month of age. Do not contaminate food containers or watering devices. Do not use where milk is produced or stored.

WARNING: Harmful if swallowed, inhaled or absorbed through the skin. Avoid spray mist or fumes. Avoid getting in eyes, on skin or on clothing. Wash thoroughly with soap and water. Change clothing immediately and wash before re-use. Keep away from children and domestic animals. Malathion, upon repeated or prolonged use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine may be used. Destroy container when empty by perforating or crushing and burying in a safe place. Do not re-use. Avoid contamination of feed or foodstuffs.

This product will kill fish. Keep out of any body of water. Apply this product only as directed on this label.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and type of application equipment will vary, Seller makes no claims other than those stated on the label.

INC. GREAT FALLS, MONTANA 59406