PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Read entire label. Follow directions carefully. Harmful if swallowed. Avoid contact with skin, eyes or clothing. In case of contact, wash with soap and water. Avoid inhalation of spray mist. Avoid contamination of feed and food stuffs. Do not store near seed, feed, insecticides, fungicides, herbicides, fertilizers. Keep out of reach of children and domestic animals. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ENVIRONMENTAL HAZARDS

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Rinse equipment and dispose of wastes by emptying in ditches and away from open water supplies. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION For many annual and perennial weeds in spring wheat, sorghum, barley, rye, oats, roadsides, pastures and turf areas. It is more effective when used on small, actively growing weeds. Use ground pressure equipment. Especially suited for boom type pressure sprayers with nozzles for spot spraying. Do not use this equipment because it sprays at a fixed rate. Proper application depends on the pressure in the system, the nozzle opening, the speed of travel and the uniformity of coverage. Poor weed control may result if pressure, speed or nozzle opening is changed. Avoid spray drift damage to sensitive crops. Do not apply when vapors from the spray may move onto susceptible crops. Do not apply when temperatures range over 90°F. Reduce spray pressure and increase the amount of water as the temperature rises. Use only enough pressure to obtain adequate coverage. Do not apply when there is danger of rain within 6 hours after application.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

APPLICATION INSTRUCTIONS

Use these recommendations only on crops listed in the Directions for Use section of this label. Do not apply this product through any type of irrigation system. Do not apply by air. Do not apply to crops other than those listed on this label.

SMALL GRAINS (WHEAT, BARLEY, RYE, OATS)

Winter Wheat and Rye Apply in the spring after the crop has begun to grow actively.

Spring Wheat, Oats and Barley Apply after the crop has reached the 3 leaf stage but before the boot stage.

FALLS CHEMICALS INC.

ACCEPTED
01-30-84
JUL 0 1984
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 40831-44

ACTIVE INGREDIENTS	
* Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	47.5%
INERT INGREDIENTS	52.5%
Total	100.0%

Contains 3.8 lbs. 2,4-D Acid per gallon

* 2,4-Dichlorophenoxyacetic Acid equivalent 39.4%
Isomer Specific by AOAC Method No. 6.D01.5

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 40831-44 EPA Est. No. 40831-MT-01

possible crop injury will be acceptable. Use higher dosage for hard to kill weeds. Treat when crop is fully tillered (4-8 inches high) up until early boot stage. NOTE: Do not spray during seedling stage, or boot, to dough stage. Do not graze or feed forage from treated fields within 2 weeks after treatment. Do not feed treated straw to livestock.

SUSCEPTIBLE WEEDS (Use lower dosage rate)

ARROWHEAD	LAMBSQUARTER	RAGWEED
BUCKHORN	MUSTARD	SUNFLOWER
COCKLEBUR	MORNING GLORY	WILD CARROT
DANDELION	PIGWEED	WILD RADISH
DOCK	PLANTAIN	WILD SWEET POTATO

HARD TO KILL WEEDS (Use higher dosage rate)

BINDWEED	HOARY CRESS	SHEEP SORREL
BUCKBRUSH	HORSE NETTLE	WILD ONION
BUCKWHEAT	LOCOWEED	WILLOW
CANADA THISTLE	MILKWEED	

CORN (Post Emergence) May be applied at several stages of growth. Use drop nozzles when corn is 10 inches or over. Mix 1/2 cup/gal in 3 gals water and wet weeds thoroughly. NOTE: Do not apply from tasseling to dough stage.

Early Spraying (Corn is 2-4 inches high) Apply as soon as possible after most weeds have emerged. Use 1 pt/acre in sufficient water to give uniform coverage. Corn drop nozzles are not necessary at this time.

Lay By Spraying (Corn is 2-3 ft. high) Use drop nozzles. At this stage of growth since stalks may become brittle from 2,4-D, there is always a chance that high winds may damage crop 1-3 days after spraying. Cultivation should be completed before applying this spray. Use 1 pt/acre.

High Clearance Spraying This spray is especially useful when weeds get started after Lay By. Use 1 pt/acre, adjust spray nozzles to hit highest weeds.

SORGHUM (Post Emergence ONLY) Use drop nozzles. Sorghums are most tolerant to 2,4-D when 4-12 inches high. Spray a few plots first. Use 1 pt/acre. NOTE: Do not apply during flowering (head) stage.

HARD TO KILL WEEDS in NON CROPLAND For perennial weeds such as Sow Thistle and Russian Knapweed, see also drainage ditchbanks, fence rows and farm roads. See Use 2-3 qts in 20 gals water with 5-8 oz liquid detergent. Treat when weeds in bloom stage. Treat in Spring and again at bloom time before weeds go to seed. Repeat treatment next year. NOTE: Do not let spray drift.

PASTURES AND TURF Use 2-4 qts in 10 gals water/acre. Apply early in Spring when weeds are actively growing. Treat again in bud stage before weeds go to seed. NOTE: Remove livestock from treated areas during spray application and for 3 days after treated areas dry.

LAWNS Use 1 qt/gal water/1000 sq ft. Repeat as needed. NOTE: Do not spray new lawns until after first mowing. Do NOT use on Bentgrass, St. Augustine grass, Dichondra or Carpetgrass lawns.

NOTICE To the extent consistent with applicable quantities and terms as indicated herein with respect to application as determined by appropriate governmental authorities and to the extent that the product is stated on the label when used in accordance with the directions under conditions of use, this warranty extends to the use of the product by purchaser or user at normal application rates under the conditions that are reasonably foreseeable by Falls Chemicals Inc. Use inconsistent with Falls Chemicals Inc. label instructions or abnormal damages resulting from the handling or application of this product. All such risks shall be assumed by buyer. FALLS CHEMICALS INC. MAKES NO WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

BEST DOCUMENT AVAILABLE

NET CONTENTS ☐ 1 ☐ 2½ ☐ 5 ☐ 30 ☐ 55 **GALLONS**

FALLS CHEMICALS INC., GREAT FALLS, MONTANA 59403