

GENERAL INFORMATION: NAVUNEX 400 LIQUID SPRAY is a selective insecticide which controls a wide variety of insect pests but does not destroy beneficial insect parasites, predators and pollinators when applied at recommended minimum rates.

Do not spray NAVUNEX 400 in conjunction with alkaline materials such as lime and lime sulfur, or with summer oils or dormant oils. NAVUNEX 400 is known to be compatible with chlorinated and phosphate insecticides. However, the selective advantage of NAVUNEX 400 LIQUID SPRAY is lost when it is used in conjunction with or alternated with pesticides which are non selective.

DIRECTIONS FOR USE: Mix the recommended dosage shown per acre in sufficient water for good coverage. Aerial Application: Approximately 1-5 gallons of water. Ground Sprayers: 10-20 gallons of water. Any conventional power operated ground sprayers or aircraft spray equipment may be used.

ALFALFA, CLOVER: Alfalfa Caterpillar: 1-2 pts. Alfalfa Webworm: 1-2 pts. Western Yellow-Striped Armyworm: 1 pt. Beet Armyworm, Variegated Cutworm: 1-2 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Make only 1 application per cutting. Allow 4 days between treatment and cutting or grazing for rates up to 1 pt./acre and a 7-day interval for rates above 1 pt./acre.

BARLEY, FLAX, OATS, WHEAT: Armyworms, Beet Webworm, Variegated Cutworm: 1-2 pts. Bertha Armyworm, Diamondback Moth: 2 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary.

Note: Do not spray within 21 days of harvest. Allow 3 days between treatment and use of treated straw as feed for dairy or meat animals.

CORN: Armyworms, Cutworms: 1-2 pts. Apply specified dosage per acre in sufficient water for complete coverage. Make one application per season when plants are 3-12 inches high and direct the spray to the lower portions of plants and to the soil around the base of the plants.

Note: Do not spray when corn is in seed, or ensilage within 28 days of treatment.

COTTON: Cotton Fleahopper: 2-3 pts. Cotton Leafworm, Darkling Ground Beetle, Western Yellow-Striped Armyworm, Beet Armyworm, Southern Garden Leafhopper: 2 pts. Black Fleahopper Complex, Cotton Leaf Perforator, Leaf Hopper, Lygus Bugs, Stink Bugs: 2-3 pts. Salt-marsh Caterpillar: 3 pts. Apply specified dosage per acre in sufficient water for complete coverage but not less than 1 gal./acre.

Note: Do not spray cotton fields within 21 days of application. Do not apply within 7 days of picking. NAVUNEX 400 may cause marginal leaf burn.

SUGAR BEETS: Beet Webworm, Variegated Cutworm: 1-2 pts. Dipterous Leaf Miners: 2 pts. Alfalfa Webworm, Beet Armyworm: 1-2 pts. Salt-marsh Caterpillar: 3 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre.

Note: Do not spray within 21 days of harvest.

SEED FIELD CROPS

ALFALFA, CLOVER: Armyworms, Lygus Bugs, Stink Bugs, Variegated Cutworm: 2-3 pts. Apply specified dosage per acre in sufficient water for complete coverage but not less than 1 gal./acre. Repeat as necessary up to 3 times during the season. Alfalfa and clover seed crop may be used for feed or forage; however, do not use for seed or forage if treated with NAVUNEX 400.



NAVUNEX 400 LIQUID SPRAY

Contains 4 Lbs. Trichlorfon Per Gallon

A Selective Insecticide for Use on Field and Vegetable Crops as Listed

ACTIVE INGREDIENTS:	
Dimethyl (2,2,2-trichloro-1-hydroxyethyl) Phosphonate	41.1%
INERT INGREDIENTS:	58.9%
Total	100.0%

WARNING: KEEP OUT OF REACH OF CHILDREN
See Side Panels for Complete Warning Statements

NET CONTENTS: ☐ 1 ☐ 5 ☐ 30 ☐ 55 GALLONS

COLORADO INTERNATIONAL CORPORATION
Commerce City, Colorado 80022

BEST AVAILABLE COPY

2/

VEGETABLES

BEANS (DRY & SNAP): Western Bean Cutworm: 1-2 pts. Armyworm, Oblique Leaf Miner, Lygus Bug, Mexican Bean Beetle, Stink Bugs, Variegated Cutworm: 2-3 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1-gal./acre.
Note: Do not apply within 14 days of harvest.

LIMA BEANS: Western Bean Cutworm: 1-2 pts. Armyworm, Oblique Leaf Miner, Lygus Bug, Mexican Bean Beetle, Stink Bugs, Variegated Cutworm: 2-3 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1-gal./acre.
Note: Do not apply more than twice per season nor within 14 days of harvest.

COWPEAS (SOUTHERN PEAS, BLACK-EYED PEAS, CROWDER PEAS): Armyworm, Oblique Leaf Miner, Imported Cabbageworm, Lygus Bug, Mexican Bean Beetle, Stink Bugs, Variegated Cutworm: 2-3 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1-gal./acre. Apply before per season.
Note: Do not apply within 14 days of harvest.

ORNAMENTALS

ORNAMENTAL FLOWERS, SHRUBS AND TREES: Armyworm, Oblique Leaf Miner, Lygus Bug, Mexican Bean Beetle, Stink Bugs, Variegated Cutworm, Fine Tip Moth, Glasswinged Plant Hopper: 2 pts. Apply specified dosage per acre using sufficient water for complete coverage but not less than 1-gal./acre. They are thoroughly wetted and must be kept wet for 2-3 hours after application. Additional applications may be required.

Note: Injury to plants may occur in certain varieties of ornamentals. Do not spray ornamentals until after plantings of these flowers have been removed for the first time, treat only a few plants before spraying the entire planting.

WARNING: May be harmful if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Always wear thoroughly wetted and saturated clothing with soap and hot water before re-use. Do not use if you get medical attention immediately.

NOTE TO FLOWERS: Do not spray before 10 a.m. or after 4 p.m. Do not spray with Aerials.

ENVIRONMENTAL CAUTION: The product is toxic to birds, fish, and aquatic life. Do not apply to water. Do not apply to areas where birds, fish, or aquatic life may be present. Do not apply to areas where birds, fish, or aquatic life may be present.

This product is toxic to birds, fish, and aquatic life. Do not apply to areas where birds, fish, or aquatic life may be present.

Do not re-use empty containers. Destroy empty containers. Do not use for any other purpose.

NOTICE: Seller warrants that this product conforms to the description of use, such as weather, temperature, and other conditions, and that Seller makes no claims other than those stated on the label.

EPA Reg. No. 4715-500

BEST AVAILABLE COPY