

Reg # 40833-4

PM-15

1972

NET WT. 40 LB.  
5 BLOCKS WEIGH 200 LB.

**SWEETLIX  
FLY CONTROL**

**RABON®  
MINERAL/VITAMIN  
MOLASSES BLOCK  
FOR CATTLE  
OR HORSES**

For control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

For control of fecal flies in manure of treated horses. Prevents development of house flies and stable flies in the manure of treated horses.

**ACTIVE INGREDIENT\***

Tetrachlorvinphos: 2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate..... 0.494%

INERT INGREDIENTS ..... 99.506%  
100.000%

\*Each lb. contains 2.24 g Rabon®  
Refers only to ingredients which are not larvicidal

**GUARANTEED ANALYSIS**

Calcium (Ca), not less than.....9.0%  
Calcium (Ca), not more than.....10.8%  
Phosphorus (P), not less than.....8.0%  
Salt (NaCl), not less than.....12.5%  
Salt (NaCl), not more than.....15.0%  
Potassium (K), not less than .....3.0%  
Magnesium (Mg), not less than.....2.0%  
Iodine (I), not less than.....0.0069%  
Selenium (Se), not less than ..... 0.00216%  
Vitamin A, min., IU/lb.....200,000  
Vitamin D-3, min., IU/lb.....50,000  
Vitamin E, min., IU/lb.....200  
Total Sugars (as invert), not less than.....12.0%

**FEED INGREDIENTS**

Cane Molasses, Soybean Hulls, Dicalcium Phosphate, Salt, Animal Fat (preserved with BHA), Potassium Chloride, Potassium Sulfate, Magnesium Sulfate, Magnesium Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Riboflavin Supplement, d-Calcium Pantothenate, Thiamine Mononitrate, Niacin Supplement, Choline Chloride, Vitamin B-12 Supplement, Folic Acid, Pyridoxine Hydrochloride, Biotin, Menadione Sodium Bisulfite Complex, Ferrous Carbonate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Copper Oxide, Copper Sulfate, Calcium Iodate, Ethylenediamine Dihydrochloride, Cobalt Carbonate, Sodium Selenite, Yeast Culture, Ethoxyquin (a preservative).

**ACCEPTED**

FEB 26 1991

U.S. DEPARTMENT OF AGRICULTURE  
Animal and Plant Health Inspection  
Center for Veterinary Medicine  
Washington, D.C. 20501  
Reg. No. 40833-4

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION  
PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment washwaters.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a dry place in original container.

**CONTAINER DISPOSAL:** Remove plastic wrap and place block in feeder. Dispose of plastic wrap in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Feed 1 block per 5 head of cattle or horses. This allows all animals equal access to blocks. Feed blocks at the rate of 0.5 oz. per 100 lb. of body weight. This intake will supply 0.07 g of larvicide per 100 lb. of body weight.
2. Place blocks where animals congregate. Locate blocks throughout the pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Cattle or horses will consume about 5 oz. per head daily. When animals consume the block at the recommended rate, it will supply 3 mg of supplemental selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE CHOICE source of salt, other minerals and vitamins.
7. SWEETLIX<sup>®</sup> Rabor<sup>®</sup> Mineral/Vitamin Molasses Block prevents the development of fecal flies in the manure of treated cattle and horses, but it is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX<sup>®</sup> Rabor<sup>®</sup> Mineral/Vitamin Molasses Block should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX<sup>®</sup> Rabor<sup>®</sup> Mineral/Vitamin Molasses Block early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX<sup>®</sup> Rabor<sup>®</sup> Mineral/Vitamin Molasses Block may be fed to animals up to slaughter and to lactating cows without withholding the milk from market either during or after treatment.

EPA Reg. No. 40833-4

EPA Est. No. 40833-IN-02

**MANUFACTURED BY**  
Pacific Molasses Company  
San Francisco, CA 94105