



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 19 1990

APR 19

300/363323
17/8OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

Mr. Dan Hickman
Pacific Molasses Company
13550 S. Indiana Avenue
Chicago, IL 60627

Dear Mr. Hickman:

Subject: Protective Clothing Statement
16:8 FREE CHOICE Mineral with R.O.L.
EPA Registration No. 40833-6
R.O.L. Rabon Molasses Block
EPA Registration No. 40833-5
Fly Control RABON Mineral/Vitamin Molasses Block
EPA Registration No. 40833-4
Your Letter Dated October 16, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

The Agency has accepted your proposed Protective Clothing Statement with the following condition:

-Change "WEAR GLOVES WHEN HANDLING" to "WEAR
CHEMICAL RESISTANT GLOVES WHEN HANDLING"

Submit five copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely,

A handwritten signature in dark ink, appearing to read "George T. LaRocca".

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

**R. O. L.
RABON[®]
MOLASSES BLOCK**

ACTIVE INGREDIENT*

0.309%

INERT INGREDIENTS	99.691%
-------------------	---------

100.000%

* Each pound contains 1.4 grams Rabon^R

****Refers only to ingredients which are not larvicidal**

FEED INGREDIENTS

Cane Molasses, Soybean Hulls, Salt, Animal Fat (preserved with BHA), Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Carbonate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Copper Oxide, Copper Sulfate, Calcium Iodate, Ethylenediamine Dihydriodide, Cobalt Carbonate, Sodium Selenite, Yeast Culture, Ethoxyquin (a preservative).

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear gloves when handling.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

CONTAINER DISPOSAL: Remove plastic wrap and place block in feeder. Dispose of plastic wrap in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Feed one block per 5 head of cattle. This allows all animals equal access to blocks. Feed blocks at the rate of 0.8 ounce per 100 pounds of body weight. This intake will supply 0.07 grams of larvicide per 100 pounds of body weight.
2. Place blocks where animals congregate. Locate blocks throughout the pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 feet minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Cattle will consume about 8.0 ounces per head daily. When animals consume the block at the recommended rate, it will supply 3.0 milligrams of supplemental selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE-CHOICE source of salt, other minerals and vitamins.
7. SWEETLIX[®] R. O. L. Rabon[®] Molasses Block prevents the development of fecal flies in the manure of treated cattle, but it is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX[®] R. O. L. Rabon[®] Molasses Block should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX[®] R. O. L. Rabon[®] Molasses Block early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX[®] R. O. L. Rabon[®] Molasses Block may be fed to cattle up to slaughter and to lactating cows without withholding milk from the market either during or after treatment.

EPA Reg. No. 40833-5

EPA Est. No. 40833-IN-02

Manufactured by
PM Pacific Molasses Company
SAN FRANCISCO, CALIFORNIA 94105

745