 <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 M St., S.W. Washington, D.C. 20460</p> <p align="center">NOTICE OF PESTICIDE Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 40833-5	Date of Issuance: OCT 10 2000
	Term of Issuance: Unconditional Registration	
	Name of Pesticide Product: Sweetix® R.O.L. Rabon® Molasses Block	
Name and Address of Registrant (include ZIP Code). PM Ag Products, Inc. 17475 Jovanna Drive Homewood, IL 60430		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		

Based on your response to the Reregistration eligibility Documents, EPA has reregistered the product listed above. This action is taken under the authority of section 4 (g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Registration under this section does not eliminate the need for control reassessment of pesticides. EPA may require submission of data at anytime to maintain the registration of your product. with regards to labeling, you must make the labeling changes listed below:

1. The chemical name of tetrachlorvinphos "2-Chloro-1-(2,4, 5-trichlorophenyl) vinyl dimethyl phosphate" should be replaced with tetrachlorvinphos: (Z)-2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate" in the ingredient statement.

2. The Precautionary Statements must be modified to read as follows:

"Hazard to Humans

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals."

The precautionary language described above should be placed directly after the First Aid section.


3. Additional user safety requirement must be added to the label directly after the item described in #2 above, and should read as follows:

"Personal Protective Equipment (PPE)

Handlers of this product must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves
- socks and shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

Signature of Approving Official: 	Date: OCT 10 2000
---	--------------------------

4. The Environmental Hazards section must be expanded to read as follows:
 "Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate."
5. Per the Agency toxicology review dated May 29, 1998 and PR Notice 2000-3 the Statement of Practical treatment section must be changed to the following title and statements:

"FIRST AID

IF SWALLOWED: - Call a Poison Control Center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told by a Poison Control Center or doctor.

**IF ON SKIN
OR CLOTHING**

- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
 - Call a poison control center or doctor for treatment advice.

In case of emergency call toll free _____.

Have the product container or label with you when calling a poison control center or doctor or going for treatment."


You may refer to PR Notice 2000-3 for guidance on placement and format of this language. A copy of the PR notice is available from the internet at the following web site: http://www.epa.gov/PR_Notices.

6. The following statements should be added the directions for use:
 "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. This product is not to be used on horses destined for slaughter."
7. This compound is toxic to fish, however, due to this formulation, it is not necessary to have the statement
 "Do not contaminate water when disposing of equipment rinse water."

A copy of your most recently submitted label is stamped "Accepted with Comments" and is enclosed.

Please submit two(2) copies of your final printed label before you release the product for shipment

Sincerely,



George LaRocca
 Product Manager 13
 Insecticide Branch
 Registration Division (7505C)

enclosure

Signature of Approving Official:	Date: OCT 10 2000
----------------------------------	-------------------

GUARANTEED ANALYSIS

Crude Protein, not less than	4.0%
Crude Fat, not less than	2.5%
Crude Fiber, not more than	13.0%
Salt (NaCl), not less than	19.0%
Salt (NaCl), not more than	22.8%
Potassium (K), not less than	2.0%
Iodine (I), not less than	0.0043%
Selenium (Se), not less than	0.00133%
Vitamin A, min., IU/lb.	20,000
Vitamin D-3, min., IU/lb.	5,000
Vitamin E, min., IU/lb.	5

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS: This product may cause skin sensitization reactions in certain individuals. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves, shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

CONTAINER DISPOSAL: Remove plastic wrap and place block in feeder. Dispose of plastic wrap in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

PM MANUFACTURED BY
ag products

INCORPORATED
Homewood, IL 60430

EPA Reg. No. 40833-5
EPA Est. No. 40833-IN-02

NET WT. 33.33 LB.
3 BLOCKS WEIGH 100 LB.



R.O.L. Rabon® MOLASSES BLOCK

For control of fecal flies in manure of treated cattle.
Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT*

Tetrachlorvinphos:2-Chloro-1-(2,4,5-trichlorophenyl)	
vinyl dimethyl phosphate	0.309%
INERT INGREDIENTS**	99.691%
	100.000%

*Each lb. contains 1.4 g. Rabon®

**Refers only to ingredients which are not larvicidal.

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT 10 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the purpose of
registration under FIFRA (7 U.S.C. 136)

40833-5

FEED INGREDIENTS

Cane Molasses, Soybean Hulls, Salt, Animal Fat (preserved with BHA), Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ferrous Carbonate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Copper Oxide, Copper Sulfate, Calcium Iodate, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Sodium Selenite, Yeast Culture, Etxoxyquin (a preservative.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Feed 1 block per 5 head of cattle. This allows all animals equal access to blocks. Feed blocks at the rate of 0.8 oz. per 100 lb. of body weight. This intake will supply 0.07g. of larvicide per 100 lb. of body weight.
2. Place blocks where animals congregate. Locate blocks throughout pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Cattle will consume about 8 oz. per head daily. When animals consume the block at the recommended rate, it will supply 3 mg. of supplemental selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE CHOICE source of salt, other minerals and vitamins.
7. SWEETLIX® R.O.L. Rabon® Molasses Block prevents the development of fecal flies in the manure of treated cattle, but it is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX® R.O.L. Rabon® Molasses block should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX® R.O.L. Rabon® Molasses Block early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX® R.O.L. Rabon® Molasses Block may be fed to animals up to slaughter and to lactating cows without withholding the milk from market either during or after treatment.



7 48936 00745 7 745