

APR 19 1989

~~300/363354~~  
17/6

Mr. Dan Hickman  
Pacific Molasses Company  
13550 S. Indiana Avenue  
Chicago, IL 60627

Dear Mr. Hickman:

Subject: Protective Clothing Statement  
16:8 FREE CHOICE Mineral with R.O.L.  
EPA Registration No. 40833-6  
R.O.L. Rabon Molasses Block  
EPA Registration No. 40833-5  
Fly Control RABON Mineral/Vitamin Molasses Block  
EPA Registration No. 40833-4  
Your Letter Dated October 16, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

The Agency has accepted your proposed Protective Clothing Statement with the following condition:

-Change "WEAR GLOVES WHEN HANDLING" to "WEAR  
CHEMICAL RESISTANT GLOVES WHEN HANDLING"

Submit five copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosures

*Hickman 4-19-89*

NET WT. 50 LB.  
22.7 Kg.



**16:8 FREE CHOICE Mineral  
with RABON<sup>®</sup> Oral Larvicide**

**Mineral and Vitamin Supplement**

For control of fecal flies in the manure of treated cattle. Prevents development of face flies, horn flies, house flies, and stable flies in the manure of treated cattle.

<b>ACTIVE DRUG INGREDIENT*</b>	
Tetrachlorvinphos: 2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate.....	0.494%
<b>INERT INGREDIENTS**</b> .....	99.506%
	100.000%

\* Each pound contains 2.24 grams RABON<sup>®</sup>  
\*\* Refers only to ingredients which are not larvicidal

**GUARANTEED ANALYSIS**

Calcium (Ca), not less than .....	15.00%
Calcium (Ca), not more than .....	17.00%
Phosphorus (P), not less than .....	8.00%
Salt (NaCl), not less than .....	14.00%
Salt (NaCl), not more than .....	16.00%
Potassium (K), not less than .....	2.00%
Magnesium (Mg), not less than .....	1.00%
Sulfur (S), not less than .....	1.00%
Iodine (I), not less than .....	0.0048%
Selenium (Se), not less than .....	0.00177%
Vitamin A, min., IU/lb .....	200,000
Vitamin D-3, min., IU/lb .....	25,000
Vitamin E, min., IU/lb .....	25
Total Sugars (as invert), not less than .....	5.00%

**FEED INGREDIENTS**

Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Potassium Chloride, Potassium Sulfate, Magnesium Sulfate, Magnesium Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Riboflavin Supplement, d-Calcium Pantothenate, Thiamine Mononitrate, Niacin Supplement, Choline Chloride, Vitamin B-12 Supplement, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Folic Acid, Biotin, Ferrous Carbonate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Copper Oxide, Copper Sulfate, Calcium Iodate, Ethylenediamine Dihydriodide, Cobalt Carbonate, Sodium Selenite, Yeast Culture, Corn Distillers Dried Grains with Solubles, Red Iron Oxide, Mineral Oil, Cane Molasses, Natural and Artificial Flavoring.

**KEEP OUT OF REACH OF CHILDREN**


SEE FREE CHOICE FEEDING DIRECTIONS ALONG WITH PRECAUTIONARY STATEMENTS ON REVERSE SIDE OF LABEL

PLANT LOCATION  
SYRACUSE, INDIANA

EPA Est. No. 40833-IN-02

ACCEPTED EPA Reg. No. 40833-6  
with COMMENTS  
in EPA Letter Dated:

Manufactured by  
Pacific Molasses Company



APR 19 1990

685

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 40833-6

**KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear gloves when handling.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not contaminate water when disposing of equipment washwaters.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a dry place in original container.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide is formulated to be fed continuously on a free choice basis to cattle to provide minerals, vitamins and Rabon<sup>®</sup> daily. Feed supplies (pasture or forage) and feeder locations should be adequate to assure mineral consumption in accordance with the intake levels below:

Cattle Weight (Pounds)	Rabon <sup>®</sup> Supplement Daily Intake		Rabon <sup>®</sup> Daily Intake (Milligrams)	Vitamin A Daily Intake (IU)
	(Ounces)	(Pounds)		
500	2.5	0.15625	350	31,250
600	3.0	0.18750	420	37,500
700	3.5	0.21875	490	43,750
800	4.0	0.25000	560	50,000
900	4.5	0.28125	630	56,250
1000	5.0	0.31250	700	62,500
1100	5.5	0.34375	770	68,750
1200	6.0	0.37500	840	75,000

If cattle consume more or less than these levels, feeders should be moved further from or closer to the general resting or watering area. If cattle still fail to consume the correct amount, discontinue feeding this free choice supplement. Feed SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide as the only FREE CHOICE source of salt, other minerals and vitamins.

**Fly Control Program**

SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX<sup>®</sup> 16:8 Free Choice Mineral with Rabon<sup>®</sup> Oral Larvicide may be fed to cattle up to slaughter and to lactating cows without withholding the milk from market either during or after treatment. Rabon<sup>®</sup> Oral Larvicide should be fed at a level to provide 70 milligrams Rabon<sup>®</sup> per 100 pounds body weight.

Care should be taken to supply a clean, fresh source of water to all animals. For additional information, please contact your SWEETLIX<sup>®</sup> representative.