



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 M St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
40833-11

Date of Issuance:
AUG 16 1999

Term of Issuance:
Unconditional Registration

Name of Pesticide Product:
VMS Pest-A-Side Block with Rabon®

**NOTICE OF PESTICIDE
Reregistration**

(Under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

**Wayne G. Bomgarden
PM Ag Products, Inc.
17475 Jovanna Drive
Homewood, IL 60430**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Documents, EPA has reregistered the subject product. This action is taken under the authority of § 4 (g) (2) (c) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Registration under this section does not eliminate the need for control reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of this product. With regards to labeling, you must make the labeling changes listed below.

1) Based on the Acute Toxicology profile for this product you must include a heading of "First Aid" or "Statements of Practical Treatment". The statements of practical treatment should be placed directly below the header "Precautionary Statements". First Aid statements should read as follows:

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF ON EYES: Flush with plenty of water. Call a physician if irritation persists."

Signature of Approving Official:

M. Douglas G. LaBrecque

Date:

16 AUG 99

2) Your product is required to have a "Note to Physicians and Veterinarians." This should be located directly below your "First Aid" Statement and should contain the following information:

- Technical Information on symptomatology;
- Use of supportive treatments to maintain life functions;
- Medicine that will counteract the specific physiological effects of the pesticide;
- Company telephone number to specific medical personnel who can provide specialized medical advice;
- Information for veterinarians.

3) The "HAZARDS TO HUMANS" statement must be modified to read as follows:

"Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

Personal Protective Equipment (PPE)


- long-sleeved shirt and long pants
- shoes and socks
- chemical resistant gloves

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

4) The header "User Safety Recommendations" must be placed in a box. The recommendations should read as follows: "Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing."

5) This compound is toxic to fish, however, due to this formulation, it is not necessary to have the statement "Do not contaminate water when disposing of equipment rinse water".

6) Under "Directions for Use", the application rate must also be expressed in amounts of active ingredient per hundred pounds body weight per month.

Signature of Approving Official: 	Date: 16 AUG 99
---	--------------------

A copy of your most recently submitted label is stamped "Accepted with Comments" and is enclosed. Please submit two (2) copies of your final printed label before you release the product for shipment.


I'll be happy to answer any questions you may have regarding this issue.

Sincerely,



George T. LaRocca
Product Manger (13)
Insecticide Branch
Registration Division 7505C

enclosure
md:a:40833-11.wpd

Signature of Approving Official: 	Date: 16 AUG 99
---	--------------------

NET WT. 33.3 LB.

15.1 Kg

VMS

INSECTICIDE • FLYING INSECTICIDE



PEST-A-SIDE BLOCK WITH RABON®

AUG 16 1999

Under the Federal Insecticide
Fungicide, and Rodenticide Act
as amended, for the pest(s)
registered under EPA Reg. N

40833-11

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT:

Tetrachlorvinphos: 2-chloro-1-(2, 4, 5-trichlorophenyl)	0.30%*
vinyl dimethyl phosphate	99.70%**
INERT INGREDIENTS**	100.00%
TOTAL	100.00%

*RABON Insecticide—Contains 1.362 grams of RABON® per pound.

**Refers only to ingredients which are not larvicidal.

EPA REG. NO. 40833-11
EPA EST. NO. 40833-AL-1

CAUTION

Keep Out of Reach of Children

GUARANTEED ANALYSIS:

Crude Protein, minimum	8.00%
Crude Fat, minimum	0.50%
Crude Fiber, maximum	7.00%
Calcium (Ca), maximum	8.50%
Calcium (Ca) minimum	7.50%
Phosphorus (P), minimum	2.00%
Salt (NaCl), maximum	12.00%
Salt (NaCl), minimum	10.00%
Magnesium (Mg), minimum	3.90%
Sulfur (S), minimum	0.25%
Cobalt (Co), minimum	0.0014%
Copper (Cu), minimum	0.009%
Iodine (I), minimum	0.006%
Manganese (Mn), minimum	0.06%
Selenium (Se)	0.0012%
Zinc (Zn), minimum	0.045%
Vitamin A, minimum	20,000 I.U. Per Lb.
Vitamin D-3, minimum	5,000 I.U. Per Lb.

FEED INGREDIENTS:

Cottonseed meal, d-hydrated alfalfa meal, dicalcium phosphate, monocalcium phosphate, salt, calcium carbonate, magnesium limestone, magnesium mica, magnesium sulfate, potassium sulfate, cobalt carbonate, cupric oxide, calcium iodate, ferrous carbonate, manganese oxide, sodium selenite, zinc oxide, vitamin A supplement, D-activated animal sterol (source of vitamin D-3), yeast culture, niacin, choline chloride, riboflavin supplement, calcium bentonite (1.7%), hemicellulose extract and cane molasses.

MANUFACTURED BY

PM ag products

INCORPORATED
1055 W. 175th Street
Homewood, IL 60430



7 48936 40093 7

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

CAUTION

This product may cause skin sensitization reactions in certain individuals. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention. Wear long-sleeved shirt and pants; chemical resistant gloves; shoes and socks for protection when handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry place in original container.

DISPOSAL: Completely empty container into application (mixer/ feeder) equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed, by State and local authorities by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Provide one block per 10-15 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1.67 lbs. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the V.M.S. Pest-A-Side Block with RABON® program but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form.

Place near watering and loafing areas. Protect from rain.

If consumption of V.M.S. Pest-A-Side Block with RABON® is above 1.67 lbs. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1.67 lbs. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding V.M.S. Pest-A-Side Block with RABON® early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall.

V.M.S. Pest-A-Side Block with RABON® may be fed to slaughter and to lactating dairy cows.

V.M.S. Pest-A-Side Block with RABON® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, V.M.S. Pest-A-Side Block with RABON® should be used in conjunction with other good management and sanitation practices.

CAUTION

Follow feeding directions. 8.9 ounces of this product will supply 3 mg Selenium. Consumption of Selenium should not exceed 3 mg daily.

TAKE TIME



OBSERVE LABEL DIRECTIONS

3/93

40093

424