# Microbiocide W

ACTIVE INGREDIENT

**INERT INGREDIENT** 

8.58 lb. per gallon.

Poly[oxyethylene (dimethyliminio) ethylene-

ACCEPTED

I UNG'OU E AND RODI

JAC MIC 104

85.0°=

ISECTIC

ICIDE /

REGIST

## KEEP OUT OF REACH OF CHILDREN

This product contains 1.29 lb. of active ingredient per gallon and weighs

# CAUTION

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID; If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL: HAZARDS: Inis pesticide is toxic to fish. Keep out of lakes, streams, of ponds. Bernits may be required for discharges containing. Mis-pesticide into lakes, streams, ponds or public water. For guidarie, contact the regional office of the Environmental Protection Agency.

### DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDE W-15 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to ita use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of 4.0 to 10.0 fluid ounces of MICROBIO-CIDE W-15 per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of MICROBIOCIDE W-15, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.0 to 10.0 fluid ounces of MICROBIO-CIDE W-15 per 1000 gallons of water (8 to 80 parts per million of MICROBIOCIDE W-15) should be employed every 2 to 5 days or as needed The frequency of addition depends upon the relative amount of bleedc.<sup>1</sup> and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE & DISPOSAL: (plastic container) Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill or bury in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:

W. S. ENGINEERING

P. O. Box 349 Ottawa, Xansas 66067

EPA EST. NO.

EPA REG. NO. 40841 -3 NET CONTENTS:

DIA BOL NO