

# ENFORCER<sup>®</sup>

## FLEA & TICK SPRAY

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Storage:** Store in a cool area away from heat or open flame.

**Dispose:** Do not reuse empty container. Wrap container and put in trash collection.

### SHAKE BEFORE USE

#### CATS & DOGS

To Kill Brown Dog Ticks and Fleas: Spray the animal from a distance of 8-12 inches. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under the body. As you spray, fluff the hair so the spray will penetrate to the skin. Make sure spray wets ticks thoroughly. Do not spray into the eyes or face. Repeat as needed.

#### PUPPIES AND KITTENS

Treat same as Cats & Dogs. Spray only on back and rub in with finger tips. Avoid treatment of nursing kittens & puppies.

#### PET SLEEPING QUARTERS

Spray pet bedding thoroughly, make sure to spray under bedding. After spray has dried, remove old bedding & replace with fresh bedding.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans

### CAUTION

Harmful if swallowed. Avoid breathing vapors. Wash hands with soap and water after use.

In the home, all food processing surfaces exposed to food and utensils should be covered during treatment or thoroughly washed before use.

Cover fish aquariums before spraying.

#### Physical or Chemical Hazards

**WARNING: FLAMMABLE** keep away from heat or open flame.

PRODUCT CODE #FTS-16

**C & J Chemical, Inc.**  
ENFORCER PRODUCTS  
P.O. Box 1068  
Cartersville, GA. 30120

# ENFORCER<sup>®</sup>

# ENFORCER<sup>®</sup>

## FLEA & TICK SPRAY FOR PETS

### KILLS FLEAS AND TICKS

Pleasant Fragrance

### REPELS BITING FLIES, GNATS, AND MOSQUITOES

**ACTIVE INGREDIENTS:** Pyrethrin 0.200%; Piperonyl butoxide, technical 0.400%; 1N-Octyl bicycloheptene dicarboximide 0.668%; 2,3,4,5-Bis(2-butylene) tetrahydro-2-furaldehyde 1.000%; Petroleum distillate 0.956%; **INERT INGREDIENTS:** 96.776%.

\*Equivalent to 0.32% (butylcarbitol(5-propylpiperonyl) ether and 0.08% related compounds.

†MGK<sup>®</sup> 264, Insecticide Synergist

\*\*MGK<sup>®</sup> Repellent 11

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

Read Label, Precautions, and First Aid Statements

NET WT. 16 oz.

EPA REG. NO. 42849-21

EPA EST. NO. 42849-GA-1

ACCEPTED

APR 2 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of registration under EPA Reg. No. 40849-21

For: 105...  
433243

NOT AVAILABLE COPY

# FUTURA

ACCEPTED  
APR 18 1985  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 433243

# FUTURA™

FLOWABLE CONCENTRATE

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Futura (*Bacillus thuringiensis*) Flowable Concentrate provides best control when caterpillars are newly hatched (1st or 2nd instar) and most susceptible to the product which must be eaten. Good leaf coverage is essential to achieve effective insect control. Higher rates are generally necessary when infestations are heavy and foliage is dense.

For armyworm control, apply to plants and thoroughly to adjacent soil when larvae are young.

To obtain a suitable mixture with water, pour the recommended amount into a ¼ filled tank and agitate. Fill tank and maintain agitation until the mixing process is complete. Futura is compatible with various insecticides, fungicides, spreaders, and stickers except those that are highly alkaline in nature. Reagitate if dilute spray is allowed to stand for more than 6 hours.

For most trees, the recommended amount of product may be applied by high pressure hydraulic sprayers in 100 gallons of water per acre. Cover foliage thoroughly, but avoid runoff. Mist blower applications may use 10-20 gallons of water per acre.

Spreaders and stickers improve coverage and are recommended.

For vine, pome, citrus and tropical fruits, use recommended amount per 100 gallons water and apply 100 to 500 gallons of spray dilution per acre.

Aerial application should use recommended rates in up to 10 gallons of water per acre during early morning or evening hours when winds are low. Spray systems which produce a 50-300 micron droplet size are preferable.

For small quantity, use 1.5 teaspoonfuls per gallon of water to provide an application rate of 21 fl. oz. / acre in 100 gallons of water.

Note: Because the active ingredient, *Bacillus thuringiensis*, is exempt from tolerance requirements, this product may be applied up to the day of harvest and in storage.

Notice: Local conditions may affect the use of Futura. Consult State Agricultural Extension or Experiment Station Specialists for specific recommendations related to local crop protection problems. Biochem Products makes no warranty, express or implied, including the warranties of commerciality, acceptable quality and/or fitness for any particular purpose concerning this material, other than those which are contained on this label.

ACTIVE INGREDIENT: *Bacillus thuringiensis* Berliner, var. kurstaki, primary powder fermentation product, potency of 12,000 International Units per milligram of product (equivalent to 54.5 billion International Units per gallon of product) ..... 15%

INERT INGREDIENTS: ..... 85%

100%

CAUTION: KEEP OUT OF REACH OF CHILDREN

**PRECAUTIONARY STATEMENTS**  
Hazards to Humans: Avoid inhalation or contact with eyes or open wounds.

**STORAGE AND DISPOSAL**  
Storage: Tightly reclose containers of unused Futura. Store in cool place. Activity may be impaired by storage at temperatures above 90°F.

Disposal: Drums can be used for non-food products after thorough cleaning with water. Smaller sized containers should not be reused. Perforate or crush and discard container according to local trash disposal regulations and in a safe place.



## Biochem Products

A Division of  
Salsbury Laboratories, Inc.  
A Member of  
the Solvay Group  
Montcharin, Delaware 19710