

MARKED-UP
COPY

ENFORCER INSECTICIDE POWDER

- Quick Kill Action and Long-Lasting Control.
- Controls Roaches, Silverfish, Ants, Bedbugs, Houseflies, Fleas, Box Elder Bugs and Crickets in and Around the Home.
- For Use in and Around the Home Including Yards, Gardens and Lawns to Control Pests on Dogs, Cats and Their Premises.
- For ~~Greenhouse~~ and Outdoor Use on Fruits, Vegetables, Flowers, Ornamentals, ~~Grains~~ and Grass up to and Including Day of Harvest.
- Kills Aphids, Whitefly, Beetles, Loopers, Mites, Leafhoppers and others as Listed on the Label.
- For Grasses, Lawns and Turf to Kill Armyworms, Caterpillars, ~~Grinch~~ Chinch Bugs, Cutworms, Fleas, Ants, and Ticks.
- Kills Fleas, Ticks and Lice on Dogs and Cats.
- Kills Fire Ants and Controls Fire Ant Mounds.
- This Container Makes _____ gallons of Fire Ant Mound Drench.
(*insert correct number based on 26 gals/#)
- Use as a Dust, in a Hose-End or Hand Sprayer.

(Note: Drawings of the Hose-End Sprayer and Pump up Hand Sprayer may be added)

ACTIVE INGREDIENTS:

Pyrethrins	0.10%
Piperonyl Butoxide, Technical*	1.00%
Diatomaceous Earth	95.00%

INERT INGREDIENTS:

.....	3.90%
TOTAL	100.00%

*Equivalent to 0.20% (butylcarbityl)(6-propylpiperonyl) ether and 0.80% related compounds.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel for Precautionary Statements

ENFORCER PRODUCTS, INC.
 P.O. BOX 1060, CARTERSVILLE, GA 30120-1060
 (800) 241-5656

EPA EST #40849-GA-1

EPA REG #40849-

ACCEPTED

Net Contents
 SEP 12 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 40849-66

274

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For Indoor/Outdoor Residential Use only

This product is a contact insecticide. Treat where insects are present and repeat as required to maintain control. ~~Apply uniformly to running areas and places where insects are found.~~ Apply uniformly with a hand duster, powder duster or similar equipment to running areas and places where insects are found.

OUTDOORS

This dust may be used on the following ~~crops~~ plants growing outdoors ~~or in greenhouses~~ up to and including the day of harvest. When applied according to use, cautions and directions, this product may be used where resistant insects are a problem. Rates vary on insect populations and foliage thickness. However, adequate control for the insects listed on this label may be obtained at the rate of 16 oz. (1 lb.) of product per 1000 sq. ft.

As a Dust: Best results are obtained when the plants are damp. Apply uniformly to insure both upper and lower leaf surfaces are covered.

In a Hose-End Sprayer: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. Add a small amount of water to the jar and add the amount of powder needed for the measured area. Stir to make slurry, then add the remaining water to the top of the jar. Empty entire contents of the jar on the premeasured area.

In a Hand Sprayer: Use 4 tablespoons (0.6 oz.) to one gallon of water. Shake container often to keep the powder agitated during application. Apply the contents to a premeasured area.

Deleted: Cotton, Cranberries, Grains, Pasture Grass, Peanuts, and Tobacco

CROP PLANT	PESTS CONTROLLED
ASPARAGUS	Asparagus Beetle
BEANS	Aphids, Leafhoppers, Whitefly, Mexican Bean Beetle, 12-Spotted Cucumber Beetle
BROCCOLI, CABBAGE, BRUSSEL SPROUTS, CAULIFLOWER	Aphids, Cross-striped Cabbageworm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug
CELERY	Green Peach Aphid, Cabbage Looper, Leaf Tier
CORN	Corn Earworm, Budworm
EGGPLANTS	Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphids
LETTUCE	Cabbage Looper, Green Peach Aphids, Diamondback Moth Larvae, Imported Cabbageworm
MUSTARD GREENS, KALE, COLLARDS, TURNIPS	Imported Cabbageworm, Diamondback Caterpillar, Aphids, Cabbage Looper
PECANS	Pecan Weevil
PEAS	(Field) Armyworms, Caterpillars, Pea Curculio
PEPPERS	Green Peach Aphids
POTATOES	Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper
PEACHES	Peach Borer
RADISHES	Aphids, Flea Beetle
SOY BEANS	Armyworms, Cabbage Looper, Mexican Bean Beetle
SPINACH	Aphids, Cabbage Looper, Webworm
TOMATOES	Green Peach Aphids, Colorado Potato Beetle, Stink Bug, Flea Beetle
ORNAMENTALS	(African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthemums, Dahlias, Dogwoods, Geraniums, Gladioli), Aphids, Flea Beetle and Leafhoppers.

GRASSES, LAWNS & TURF

To Kill Armyworms, Caterpillars, ~~Grass~~ Chinch Bugs, Cutworms, Fleas, Ants, And Ticks: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. ~~Apply evenly using a Hose-End Sprayer.~~ Apply either as a Dust or as a Wettable Powder.

As a Dust: The best results are obtained when applied with a hand or power duster or other suitable means of application to the infested area. Apply lightly and uniformly to the treatment area.

In a Hose-End Sprayer: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. Add a small amount of water to the jar and add the amount of powder needed for the measured area. Stir to make slurry, then add the remaining water to the top of the jar. Empty entire contents of the jar on the premeasured area.

In a Hand Sprayer: Use 4 tablespoons (0.6 oz.) to one gallon of water. Shake container often to keep the powder agitated during application. Apply the contents to a premeasured area.

FIRE ANTS

To Kill Fire Ants and Keep Them from Entering Homes: Thoroughly dust patios, barbecue areas, driveways, entry ways, door sills, window frames, other entrance areas, foundations of buildings, ant trails, around pools and areas where ants are noticed.

Fire Ant Mounds: For best results, dust the perimeter of the mound first. With a stick, disturb the mound, then dust mound in a circular motion working toward the center. Dust until the area is well covered, contact as many insects as possible.

For Use as a Mound Drench: An ant mound 12-14 inches diameter requires approximately one gallon of the mix. Prepare solution by mixing 4 tablespoons (0.6 oz.) per gallon of water. Shake well immediately prior to use. Drench both the mound and the area within six inches of the mound with solution. Apply gently to avoid disturbing ants. Reapply as necessary. For best results, apply in cool weather (65°F to 80°F) or in early morning or late evening hours. Treat new mounds as they appear.

ANIMAL QUARTERS

(including, but not limited to Barns, Pet Kennels, Runs and Stables).

To Control Gnats, Flies, Mosquitoes, Lice, Mites, Fleas and Ticks: Use at the rate of 2 oz. per 100 sq. ft. of surface area. Apply to the hiding places where the pests are found. Repeat the treatment as necessary to maintain control.

INDOORS

Remove or cover food prior to use. All food preparation surfaces and utensils should be covered during treatment, or thoroughly washed before use.

~~Use at the rate of 2 oz. per 100 sq. ft. of surface area.~~ Use a bulbous and/or powder duster to apply as a spot/crack and crevice treatment indoors.

Roaches and Silverfish: Dust lightly, paying special attention to cracks and hidden surfaces, around pipes, sinks and moldings.

Houseflies: Lightly coat window frames and sills so flies lighting will contact dust.

Ants, Crickets and Box Elder Bugs: Dust lightly areas around windows, doors and other places where pests may enter premises.

Bedbugs: Treat liberally baseboards, moldings, cracks, bedstead and both sides of mattresses.

DOGS AND CATS

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS AND CATS**

Fleas, Ticks and Lice: Lightly work the powder into the hair of the pet. Use brush/comb to work material down to the skin. Pay strict attention to places where the animal cannot reach, such as the scalp. Dust ears of the pet with care. Individual ticks observed should be dusted thoroughly. Dust the pet's bedding and areas where the pet sleeps, as well as cracks and crevices adjacent to them. Repeat once a month or as necessary to maintain control.

Do not use on pets under 12 weeks old. Consult a veterinarian before using this product on debilitated, aged, pregnant, nursing or medicated animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal

STORAGE: Store in cool, dry place.

DISPOSAL: Do not reuse container. Wrap and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Avoid contact with eyes and skin. Use adequate ventilation and avoid breathing dust. Use a dust mask for prolonged periods of use. Wash hands thoroughly after using and before smoking or eating. Cover fish aquariums before dusting.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting

IF IN EYES: Flush with plenty of water. Call a physician immediately if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF INHALED: Remove the person to fresh air.

ENVIRONMENTAL HAZARDS

Do not apply when weather conditions favor drift from treatment area. This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters.

This container is sold by weight, NOT by volume. You can be assured of the proper weight even though some settling of contents normally occurs during transportation and handling.