

(Contains 5.2 pounds of SBP-1382 per gallon)

For indoor use in ULV, Mechanical Misting Sprayers, or Thermal Foggers

For use in food processing plants, milk handling areas of milk, cheese and ice cream plants, bakeries, breweries, bottling plants, grain mills, warehouses, boxcars, stores, institutions, homes, and horse stables, not for application if horses will be used for food.

Active Ingredients:

*† ● resmethrin ((5-(phenylmethyl)-3-furanyl) methyl 2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropane-carboxylate	3.00% w/w
Inert Ingredients:	97.00%
	100.00%

*Cis/traps Isomers ratio: max. 30% (±) cis and min. 70% (±) trans.

†U.S. Patent Nos. 3,465,007 & 3,542,928

● Penick's SBP-1382® brand of resmethrin insecticide, Penick Corp., 1050 Wall St. West, Lyndhurst, N.J. 07071

MAY 6 1987

CAUTION: KEEP OUT OF REACH OF CHILDREN
See side panel for additional precautionary statements.

Net Contents: 55 gallons

PA Reg. No. 40871-4
PA Est. No. 40871-CA-1



DOT SHIPPING NAME
INSECTICIDE
LIQUID
N.O.S.
COMBUSTIBLE
NA 1993

MANUFACTURED BY

Extinguish all flames and all air conditioning and ventilating equipment. Close all doors and windows after spraying. Ventilate room before reentry.

SPACE SPRAY: To kill adult and accessible insects: grain weevils, Flour Beetles, Saw-toothed Grain Beetles, Mediterranean Flour Moths, Indian Meal Moths, Spider Beetles. Direct output into corners, under packaged goods hitting as many of the insects as possible and spray upward toward upper corners of room and ceiling. To kill ants and gnats, treat as a space spray directing output toward upper corners and ceiling.

CONTACT SPRAY: Use undiluted to kill cockroaches, silverfish, earwigs, and spiders, crickets, centipedes, sowbugs, millipedes, carpet beetles. Apply the spray as a coarse wet spray into all cracks and crevices in woodwork, walls, floors beneath sinks and other insect harborages, contacting as many insects as possible.

FOR MECHANICAL MISTING EQUIPMENT SUCH AS THE MICROSOLO OR CHALLENGER MACHINE.

Misting instructions:
Dilute to 0.5% SURESECT-300 by adding 1 fl. oz. of the 3.0% concentrate to 5 fl. oz. of petroleum distillate.
Dilute to 0.25% SURESECT-300 by adding 1 fl. oz. of the 3.0% concentrate to 11 fl. oz. of petroleum distillate.
Dilute to 0.20 SURESECT-300 by adding 1 fl. oz. of the 3.0% concentrate to 14 fl. oz. of petroleum distillate.
Dilute to 0.10% SURESECT-300 by adding 1 fl. oz. of the 3.0% concentrate to 29 fl. oz. of petroleum distillate (which would be used for flying insects only). For the most effective results against flying insects, operate spraying dispenser adjusted to deliver a fine mist. For flying insects, use either 0.10% or 0.25% SURESECT-300.

To kill flies, mosquitoes, gnats, and small flying moths: Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room before reentry. Repeat as necessary. To kill crawling insects, adjust spray to produce large wetting type spray particles and use either 0.25% or 0.50% SURESECT-300. To kill cockroaches, earwigs, grain mites, sowbugs, spiders, silverfish, crickets, centipedes and gain insects (adults and accessible stages of confused flour beetles, red flour beetles, saw-toothed grain beetles, shiny spider beetles, carpet beetles, grain weevils, Angoumois grain moths, Indian meal moths): Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, storage areas and other insect harborage areas. Spray directly on insects where possible. Repeat as necessary. To kill ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

THERMAL APPLICATIONS, INDOORS: Use 0.2% SURESECT-300 for the control of flies, mosquitoes and gnats in warehouses, restaurants and industrial institutions. Shut off ventilating systems, air ducts, air-conditioning equipment and close off room or area to be treated. Extinguish all flames and pilot lights. Apply as a dry fog at the rate of 1-1/3 fluid ounces per 1000 cubic feet of space. Direct nozzle of thermal applicator toward the upper corners of the room and along ceiling. Close off room for at least one-half hour. Do not remain in treated areas. Ventilate the area before reentry.

WARRANTY STATEMENT: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

BEST AVAILABLE COPY

... while food processing is under way. After
... these areas are ventilated to remove insecticide
... facilities and equipment are thoroughly washed with
... detergent solution and rinsed with potable water to
... traces of contamination. In the home, all food processing
... and utensils should be covered during treatment and
... washed with an acceptable detergent and rinsed with

STORAGE AND DISPOSAL

PRODUCT STORAGE: Keep away from heat. Store in dry area. Keep container lid closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

This concentrate is designed for use undiluted in ULV sprayers such as the LECO, ULV or Micro-Gen, which give particles of aerosol size. It can also be diluted with oils which meet Food Additive Regulation 21.1182 for use in mechanical fogging equipment such as the Microsol or Challenger machines which give larger particles. It can also be diluted with oil as specified and applied as a contact spray.

In food handling areas, food should be removed or covered previous to treatment. All food processing surfaces should be covered or thoroughly cleaned before using. Apply only when facility is not in operation.

Before application, remove plants and animals from area to be treated. Remove horses before spraying stables.

E
E

M