


<p>US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20480</p> <p>NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i></p>	<p>EPA REGISTRATION NO. 40871-9</p>	<p>DATE OF ISSUANCE 06/04/89</p>
<p>TERM OF ISSUANCE</p>		
<p>NAME OF PESTICIDE PRODUCT EXTRACT-7</p>		
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP code)</p> <p>Meridian, Inc. P.O. Box 100 Durham, NC 27709-0100</p>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>In addition, your responsibility to the General Registration Standard - "Guidance for the Registration and Reregistration of Pesticide Products Containing Polymers of Calcium Alginate" as the Sole Active Ingredient"--issued February 1986 and revised April 1986, EPA has reregistered the product listed above in accordance with FIFRA section 6(d). EPA has also reviewed your response to the Pesticide Label Improvement Program issued July 25, 1987 and has determined that your product is in compliance with the program.</p> <p>Your amended label has been stamped approved and a copy is enclosed. You must incorporate my comments noted on your label. This label must be on products released for shipment within 1 year or by the next label printing, whichever occurs first. This label must also appear on all supplemental registrations within 1 year of this Notice of Reregistration or at the next label printing, whichever occurs first.</p> <p>Your product meets the criteria for reregistration under this Standard. Failure to comply with this Standard may result in the Agency's issuance of a Notice of Intent to Cancel the registration under FIFRA section 6(t)(1).</p> <p style="text-align: right;">  Jeff Kempter Product Manager (32) Antimicrobial Program Branch Registration Division (75-767C) </p> <p><input type="checkbox"/> Enclosure <input type="checkbox"/> ATTACHMENT IS APPLICABLE</p> <p>SIGNATURE OF APPROVING OFFICIAL _____ DATE _____</p>		

EXTRACT-2

ACTIVE INGREDIENT
SODIUM
HYPOCHLORITE 12.5%

INERT
INGREDIENTS 87.5%
with COMMENTS
in EPA Letter Dated

JUN - 2 1989

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

40871-9

**KEEP
OUT OF
REACH
OF CHILDREN**

DANGER

FIRST AID: If on skin,
wash with plenty of soap
and water. If in eyes, flush
with water for at least 15 min-
utes. Get medical attention. If
swallowed, drink large quantities of
milk, or gelatin solution or, if these are
not available, drink large quantities of
water. Do NOT give vinegar or other acids.
Do NOT induce vomiting. Get prompt medical
attention

See additional precautions on side panel

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. May cause severe skin irritation or chemical burns to broken skin.
Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves
when handling this product. Wash after handling. Avoid breathing vapors. Use in poorly ventilated areas as
soon as possible. Do not return until odors have dissipated.


ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes,
streams or ponds. Treated effluent may not be discharged into lakes, streams, ponds or public waters without
a valid discharge permit. For guidance, contact the regional office of the Environmental Protection Agency.

PHYSICAL AND CHEMICAL HAZARDS: STRONG

OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product
with grass seed, lime, urea, etc. or with ammonium acids, detergents or other chemicals may release
hazardous gases irritating to eyes, lungs and mucous membranes.

STORAGE AND DISPOSAL: Store in a cool, dry area away from direct sunlight.
Keeps 12 months. Avoid areas with large quantities of water. Rinse empty container thoroughly with water and
return to manufacturer or dealer by placing in trash collection or burying in an approved landfill.
Product or residue that cannot be used should be diluted with water and disposed of in a sanitary manner. Do
not contaminate food or feed by storage, disposal or cleaning of equipment.

DOT
SHIPPING NAME
COMPOUND
CLEANING LIQUID
NA 1760



manufactured by
CHEMLAND, INC.

1298 S WALNUT
TURLOCK, CA 95381

1792 LATHAM ST
MEMPHIS, TN 38116

EPA REG. NO. 40871-9
EPA EST. NO. 40871-CA-1

NET CONTENTS:

DIRECTIONS FOR USE GENERAL CLASSIFICATION

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT
IN A MANNER INCONSISTENT WITH ITS LABELING

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

SANITIZATION OF NONPOROUS FOOD CONTACT SURFACES

RINSE METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

IMMERSION METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

CLEAN-IN-PLACE METHOD - Thoroughly clean equipment after use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 2 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under pressure for at least 10 minutes to insure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine. Rinse system with potable water prior to use.

SPRAY/FOG METHOD - Pre-clean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 2 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 6 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with a 600 ppm solution with a 200 ppm solution.

AGRICULTURAL USES

FOOD BCG SANITIZATION - Thoroughly clean all eggs. Thoroughly wash product with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 110°F. Immerse sanitizer so that the eggs are thoroughly wetted. Allow to stand for 2 minutes. Allow to dry before casing or breaking. Do not apply a potable water wash. Should not be re-used to sanitize eggs.

FRUIT & VEGETABLE WASHING - Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 5 oz. of this product in 200 gallons of water to produce a sanitizing solution of 25 ppm available chlorine. After sanitizing fruit or vegetables for 2 minutes in a second wash tank, immerse in a recirculating sanitizing solution. Spray rinse vegetable surfaces with a sanitizing solution prior to packaging. Rinse fruit with potable water before packaging.

LAUNDRY SANITIZERS

Commercial Laundry Sanitizers

Wet fabrics or clothes should be spun dry prior to sanitizing. Mix 2 oz. of this product with 10 gallons of water to yield a 200 ppm available chlorine solution. Promptly after mixing the sanitizer, add the detergent. Pre-wash prior to washing fabrics/clothes in the regular detergent. Test the level of available chlorine, if solution is low, add more to stand. Add more of this product if the available chlorine has dropped below 200 ppm.

FARM PREMISES

Remove all animals, poultry, and feed from premises, wash and disinfect. Remove all litter and manure from floors, walls and surfaces of stalls, chutes and other facilities occupied or traversed by animals and poultry. Empty all troughs, racks and other feeding and watering equipment. Thoroughly clean all surfaces with soap or detergent and rinse. Disinfect, saturate all surfaces with a solution of at least 100 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution may be used for thorough mixing 11 oz. of this product with 10 gallons of water. Clean halters, ropes and other types of equipment used in handling animals or poultry, as well as the cleaned floors, aisles, and areas. Remove litter and manure. Ventilate buildings, care, and equipment. Do not house livestock or poultry or employ equipment until the available chlorine has been dissipated. All treated feed racks, mangers, troughs, feeders, fountains and waterers must be rinsed with potable water.

SWIMMING POOL WATER DISINFECTION

For a new pool or spring start-up, superchlorinate with 10 times the amount of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine. Check the level of available chlorine with a test kit. Adjust water pH to between 7.2 to 7.6. Adjust and maintain the available chlorine residual between 50 to 100 ppm.

To maintain the pool, add manually or by a feeder device 1.0 ppm available chlorine for each 10,000 gallons of water to yield an available chlorine residual of 0.6 to 1.0 ppm by weight. Stabilized pools should maintain a residual of 1.0 to 1.5 ppm available chlorine. Test the pH, available chlorine, and alkalinity of the water frequently with appropriate test kits. Winter treatment will depend upon temperature and number of swimmers.

Every 7 days, or as necessary, superchlorinate the pool with 10 times the amount of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Maintain the pool until the chlorine residual is between 1.0 to 3.0 ppm.

At the end of the swimming pool season or when water is to be drained from the pool, chlorine must be allowed to dissipate from treated water before discharge. Do not chlorinate the pool within 24 hours prior to discharge.

WINTERIZING POOLS - While water is still clear & clean, add 1.0 ppm available chlorine per 1000 gallons, while filter is running, to obtain a 3.0 ppm residual, as determined by a suitable test kit. Cover pool and heater components for winter by following manufacturer's instructions.