

FROM
PYROCIDE INTERMEDIATE 5192

ACTIVE INGREDIENTS:

Pyrethrins	0.30%
*Piperonyl butoxide, technical	0.60%
±N-Octyl bicycloheptene dicarboximide	1.00%
INERT INGREDIENTS:	98.10%
TOTAL 100.00%	

*Equivalent to .48% (butylcarbityl) (6-propiperonyl) ether and .12% related compounds.
±MGK 264, Insecticide Synergist.

ONLY FOR SALE TO, USE AND STORAGE BY SERVICE PERSONS

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional
precautionary statements.

EPA REG. NO. 40871-25

It is a violation of the law to use this product in a manner inconsistent with its label.

To protect dairy cows and cattle from flies, mosquitoes and gnats: Spray shortly before morning or evening milking. Apply as a light mist (2) ounces per animal. Apply from a distance of 10 to 15 feet. Direct the spray particles to collect upon the hair coat. Pay particular attention to shoulder blades, belly and legs. Wash teats and udders with soap and warm water prior to milking. Do not soak the hide.

For Fogging: Use approximately one (1) ounce of material for each 1,000 cubic feet of space. Keep doors and windows closed for at least ten minutes. Sweep up and destroy all fallen flies. Follow recommendations of equipment manufacturer for proper positioning of the fogging machine.

For Calves: Spray very lightly once or twice daily.

Other Livestock, Horses and Swine: Spray very lightly once or twice daily.

To Protect Cattle and Other Livestock (see above) from Attack by Lice and Ticks: The animals should be sprayed at the rate of two (2) ounces per spraying for adult animals the size of a cow. Immediately rub the spray into the hair, paying particular attention to infested areas. Repeat as necessary, but not more frequently than once every three weeks. Ticks must be contacted directly with the spray.

To Protect Cattle and Other Livestock (see above) from Attack by Fleas: Spray bedding and walls of pen, particularly cracks and crevices. The livestock should be sprayed daily until infestation is controlled.

To Protect Cattle and Other Livestock (see above) from Attack by Face Flies: Apply spray to the faces of cattle at the rate of one (1) ounce per animal, paying particular attention to the areas near the eyes and nostrils. Wipe on with a heavy cloth or sponge, or spray on with a sprayer that delivers coarse droplets. Avoid spraying directly into the animal's eyes. Treatment should be made daily while the cattle are stanchioned for the morning milking to give protection from face flies.

NET CONTENTS _____ GALLONS

CHEMLAND

Manufactured by
CHEMLAND, INC.

**DOT
SHIPPING NAME**

INSECTICIDE
LIQUID
N.O.S.
COMBUSTIBLE
NA 1993

BEST AVAILABLE COPY

MAY 6 1987

U.S. Environmental Protection Agency
Pesticide and Fertilizer Act
is authorized for the pesticide
registered under
EPA Reg. No. 40871-25

STORAGE AND DISPOSAL

PRODUCT STORAGE: Keep away from heat. Store in dry area. Keep container lid closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.