


ACCEPTED
OCT 1 1978
40926-2

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



67-33

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:	By Wt.
Methyl bromide	67%
Chloropicrin	33%
Total	100%

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)

DANGER
KEEP OUT OF REACH OF CHILDREN
POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNINGS ON SIDE PANEL

NET WEIGHT _____ LBS.

EPA Reg. No 409262

LOT NO _____

HyYield-Bromine

P.O. Box 786

EL PASO, N.M. 79905

Many pesticides are poisonous and may leave a toxic residue on the plants to which they are applied. The U. S. Food and Drug Administration has established maximum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of these amounts. The "Fumigation Directions" are based on the best available information, and if followed carefully should not leave excessive residues at harvest. However, HyYield-Bromine assumes no responsibility as to their accuracy nor for any loss due to excessive residues.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the use of it at such use.

Empty cylinders should be returned to HyYield-Bromine location from which the equipment was made. Consult local sales office if questions arise.

40926-2

67-33 FUMIGATION DIRECTIONS RECOMMENDED APPLICATION RATES

Soil treatment for
Strawberries 225 lb. 325 lb. Acre

Expose to fumigation for 14 days. Aerate 2 weeks before setting the plants in treated areas

Soil treatment for
Tomatoes 225 lb. 325 lb. Acre

Expose to fumigation for 14 days. Aerate 2 weeks before setting the plants in treated areas

CITRUS SOIL FOR CONTROL OF PHYTOPHTHORA IN FLORIDA SANDY SOILS
This is a preplant or preplant treatment. Trees which are planted in the treated soil should not have harvestable fruit for at least 24 months. Use a minimum of 225 lb. 325 lb. 100 sq. ft. Expose to fumigation for 14 days. Covering treated area with a 4 ft. tarpaulin. Will control disease for at least 4 years. Aerate 2 weeks before setting the plants in treated area

Apply with a soil applicator or by hand to a depth of 8 inches. Use a 4 ft. tarpaulin drag or similar device to cover the treated area. Cover with a 4 ft. tarpaulin. Remove tarpaulin after 14 days. Do not get on bare soil for at least 2 weeks before setting the plants in treated area

Do not apply to soil which is wet or has a high water table

Do not apply to soil which is wet or has a high water table

Do not apply to soil which is wet or has a high water table

ACCEPTED

DEC 1 1978

U.S. DEPT. OF AGRICULTURE
FEDERAL EXTENSION ACT
Approved for the pesticide
Federal Order
40926-7

**POISON
DANGER**

**MAY BURN SKIN AND EYES
POISONOUS IF INHALED**

**Do Not Breathe Vapor
Do Not Get in Eyes, on Skin or Clothing**

In case of contact immediately flush skin or eyes with large amounts of water for at least 15 minutes and get medical attention for eye contact. If liquid is spilled on clothing or shoes, remove them immediately and air thoroughly. Do not reuse shoes or clothing until free of all contamination.

POISON

**SEND FOR DOCTOR IN CASE OF POISONING
ANTIDOTE:** Re-examine patient to free air passages, give artificial respiration, use antidote recommended by treating physician. Do not induce vomiting. Do not give anything by mouth until advised by a physician.

NOTICE

Following soil fumigation, undesirable concentrations of chloropicrin may drift to nearby areas. If this occurs, immediately cover treated area with a plastic tarpaulin. Leave tarpaulin on overnight and remove during the daytime. Recover the area if eye irritation indicates that the undesirable concentration is continuing. These concentrations are most likely to occur in the evening or at night when the air is most likely to be still. Do not make application of any substance there is little or no air movement or there is an inversion condition in the atmosphere.

NOTE CAREFULLY: Following fumigation the level of soluble salts and ammonia nitrogen may be raised. This is most likely to occur when heavy rates of fumigant and fertilizers are applied to soils that are either wet or contain high organic matter. Fertilizers containing ammonium salts should not be used. Apply only fertilizers containing nitrogen after the soil has been established and the soil is free of the fumigant. For more information on the use of fertilizers, consult the manufacturer's literature. Do not apply fertilizers to soil which is wet or has a high water table. Do not apply fertilizers to soil which is wet or has a high water table.