

# Vet Shock

### PELIGRO

#### WARNING

#### NOTE TO PHYSICIANS

#### DO NOT STORE BELOW FREEZING TEMPERATURE DIRECTIONS FOR USE

- REQUIRED: TECHNICAL EQUIPMENT: (1) 1/2" AT 1/2"
- All applications should be made in the early morning or late afternoon, when the temperature is below 85°F and the humidity is below 80%. Do not apply to plants that are stressed or have been treated with other insecticides. Do not apply to plants that are in bloom or have young leaves.
1. Mix the concentrate with water in a ratio of 1:100.
  2. Apply to the undersides of leaves.
  3. Repeat applications as needed.

Manufactured by  
**LEXTRON INC.**  
Greenwood, Mississippi

OCT 1981

EPA #

176740

## CYGON<sup>®</sup> 2-E Systemic Insecticide

### CONTROLS HOUSEFLIES AND INSECTS AND MITES ON ORNAMENTAL PLANTS

### WARNING! KEEP OUT OF REACH OF CHILDREN See side panel for antidote and other warnings

Lea el panel lateral con respecto al antidoto  
y otras precauciones

Active ingredients:  
Cyfluthrin (10%), diazinon (10%), methoxyacrylamide (10%),  
and pyrethrin (10%)  
Inert ingredients

EPA Reg. No. 40-942-H

NET CONTENTS: 1 GALLON

Apply to the undersides of leaves and to the soil around the plants.

Residual Warning:  
For the control of houseflies, including poultry houses and other farm buildings, apply to the undersides of leaves and to the soil around the plants. Repeat applications as needed. Do not apply to plants that are in bloom or have young leaves.

For the control of insects and mites on ornamental plants, apply to the undersides of leaves and to the soil around the plants. Repeat applications as needed. Do not apply to plants that are in bloom or have young leaves.

Do not apply to plants that are in bloom or have young leaves. Do not apply to plants that are stressed or have been treated with other insecticides. Do not apply to plants that are in bloom or have young leaves.

For more information, contact your local extension office.

#### OUTDOOR USE

CYGON 2-E is generally effective in controlling houseflies, including poultry houses and other farm buildings, and insects and mites on ornamental plants. Make adequate sprays when observed. Do not overdose or overuse. Do not use on ornamental plants grown as a source of food for livestock or on plants containing oil. Do not use on plants listed below are to

