

DEC 13 1994

Michael M. Watts  
Kincaid Enterprises, Inc.  
Box 549  
Nitro, WV 25143

Dear Mr. Watts:

Subject: Deleted uses  
Marlate 50 Insecticide Wettable Powder  
EPA Registration No. 41014-2  
Amendment application of April 13, 1994

The amendment referred to above, submitted in connection with registration under Federal Insecticide, Fungicide and Rodenticide Act, is acceptable provided that you:

Submit five (5) copies of your final printed labeling before you release the product for shipment.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6 (e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamp copy of the labeling is enclosed for your records.

Sincerely yours,

Dennis H. Edwards, Jr.  
Product Manager (19)  
Insecticide Rodenticide Branch  
Registration Division (7505C)

Enclosure



# MARLATE 50

## INSECTICIDE

### WETTABLE POWDER

#### For Dairy And Beef Cattle & Certain Other Livestock

#### Crops • Buildings • Ornamentals

#### ACTIVE INGREDIENT:

\*Methoxychlor, Technical ..... 50%

INERT INGREDIENTS ..... 50%

\*50% Methoxychlor, Technical is equivalent to 44% 1,1,1-Trichloro-2,2-bis(4-methoxyphenyl)ethane and 6% impurities associated with the active ingredient.

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat.

If on skin: Wash thoroughly with soap and water.

If in Eyes: Flush with plenty of water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

DEC 13 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

41014-2



## NET 4 LBS.

EPA REG. NO. 41014-2

EPA Est. No. 4-JM-TM-1

# KINCAID ENTERPRISES, INC.

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS &  
DOMESTIC ANIMALS

Precautionary Statements: Caution - Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Reclose container after each use. Store in a cool, dry place away from food or feed. If spilled, sweep up and dispose of below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag, then dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of the smoke.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### FARM BUILDINGS (Except Poultry Houses)

For control of Stable Fly, Housefly, Face Fly, Fleas, Mosquitoes: Use 1 lb in 2½ gal water. Apply 1 gal of spray to 500 sq ft of surface, treating walls, floors, cracks, crevices, baseboards, stalls, and pens. Such heavy deposits usually are effective for 3 to 6 weeks unless removed by weathering or other means. Care should be taken to avoid contamination of milk handling equipment, feed-stuffs, feed troughs and water receptacles. Dairy animals should not be present while spraying. The use of any residual fly spray should be supplemented with proper sanitation measures, including spraying or removal of breeding sources such as manure in and around buildings.

### STORED SEED - SEED POTATOES

Potato Tubeworm: Sprinkle MARLATE 50 over potatoes at 0.2 oz per 100 lbs before placing in storage. Do not use treated seed for food or feed purposes.

### FLOWERS, ORNAMENTALS

For control of Blister Beetles, Cankerworms, Cucumber Beetles, Flea Beetles, Fleahoppers, Flower Thrips, Japanese Beetle, Leafhoppers, Rose Chafer, Rose Slugs (Sawflies): Use 2 tablespoonfuls per gal; spray thoroughly to runoff. Begin applications when insects first appear and repeat at 7 to 14-day intervals or as needed.

### SHADE TREES AND SHRUBS

For control of Tent Caterpillars, Cankerworms, Japanese Beetle: Use 2 tablespoonfuls per gal; spray thoroughly to runoff. Begin application when insects first appear and repeat at 7 to 14-day intervals as needed.

### LIVESTOCK - DAIRY CATTLE

Direct dusting for control of Horn Fly: use 1 rounded tablespoonfuls per animal. Sprinkle on poll, neck (top and sides) and back and upper sides; rub in lightly to carry powder beneath the hair. Repeat treatment every 3 weeks as needed. Do not use on dairy goats. Do not apply to lactating animals within five (5) hours of milking or during milking.

### LIVESTOCK - BEEF CATTLE, SHEEP, SWINE, NON-LACTATING DAIRY CATTLE, NON-LACTATING GOATS

For control of Horn Fly, Lice and Fleas: Repeat treatment every 3 weeks as needed; may be used on young dairy animals up to 2 weeks of freshening. Spray or Dip - Use 2 lbs in 25 gal water. As a spray, wet animals to the skin. For control of Tail Louse of cattle, use 4 to 6 lbs in 25 gal water and apply as a spray on the infested tail only. Do not dip calves under 1 month of age. Direct Dusting - Dust animal thoroughly; rub in lightly to carry powder beneath the hair. Application on sheep, following shearing, gives temporary relief from Sheep Ticks (Keds).

## GARDEN AND ORCHARD

Use 2 tablespoonfuls MARLATE 50 per gal water and apply as a thorough spray. MARLATE 50 may also be used as a dust by diluting with any suitable diluent, preferably talc or pyrophyllite. Begin application at first sign of infestation (for fruits, begin at petal fall) and repeat at 7 to 14-day intervals or as needed. Do not apply to crops within the number of days specified before harvest or grazing as shown by PHL (preharvest limitation).

### FRUITS

**APPLE, PEAR, QUINCE** - For control of Apple Maggot, Codling Moth, Japanese Beetle, Plum Curculio, Tent Caterpillars. PHL: 7 days.

**APRICOT, CHERRY, NECTARINE, PEACH, PLUM, PRUNE** - For control of Cherry Fruitworm, Cherry Fruit Flies, Japanese Beetle, Plum Curculio, Rose Chafer, Tent Caterpillars, Cankerworms. PHL: 7 days for cherries, plums, prunes; 21 days for apricots, nectarines, peaches.

**BLUEBERRY, CURRANT, GOOSEBERRY** - For control of Japanese Beetle, Cranberry Fruitworm, Leafhoppers, San Jose Scale (Crawlers), Strawberry Rootworm, Blueberry Flea Beetle Larvae, Blueberry Leaf Beetle, Blueberry Leaf Tier Larvae. PHL: 14 days.

**BLACKBERRY, LOGANBERRY, RASPBERRY, BOYSENBERRY, DEWBERRY, YOUNGBERRY, STRAWBERRY** - For control of Rose Chafer, Strawberry Weevil, Flea Beetles, Omnivorous Leaf Tier, Spittlebugs, Japanese Beetle, Rednecked Cane Borer. PHL: 14 days.

**GRAPE** - For control of Berry Moth, Grape Leafhopper, Japanese Beetle, Leaf Skeletonizer, Rose Chafer. PHL: 14 days.

### VEGETABLES

**ASPARAGUS** - For control of Asparagus Beetle. PHL: if applied within 3 days of harvest, remove residues by washing or blanching.

**BEANS, BLACK-EYED PEAS** - For control of Corn Earworm, Cucumber Beetles, Fall Armyworm, Alfalfa Webworm, Garden Webworm, Flea Beetles, Mexican Bean Beetle, Potato Leafhopper, Bean Leaf Beetle, Japanese Beetle. PHL: 7 days; can be applied within 3 days of harvest if vines are not used for feed or forage.

**BRUSSELS SPROUTS, BEETS, BROCCOLI, CABBAGE, CARROT, CAULIFLOWER, COLLARDS, EGGPLANT, KALE, KOHLRABI, LETTUCE, PEPPER, RADISH, RUTABAGA, SPINACH, TURNIP** - For control of Blister Beetles, Flea Beetles, Leafhoppers, Alfalfa Looper, Fall Armyworm, Japanese Beetle, Imported Cabbageworm. PHL: 14 days, except: cauliflower eggplant, kohlrabi, pepper, and rutabaga, 7 days; cabbage, 3 days.

**SWEET CORN** - For control of Fall Armyworm, Armyworm, Japanese Beetle, Flea Beetles. PHL: 7 days.

**CUCUMBERS, MELONS, PUMPKIN, SQUASH, SUMMER SQUASH** - For control of Squash Vine Borer, Fall Armyworm, Cucumber Beetles, Flea Beetles. PHL: 7 days.

PEAS - For control of Alfalfa Looper, Pea Weevil (to foliage early bloom before eggs are laid and again 5 days later), Bean Leaf Beetle. PHL: 7 days.

POTATO (IRISH) - For control of Colorado Potato Beetle, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetles. PHL: None.

SWEET POTATO, YAM - For control of Fall Armyworm, Flea Beetles. PHL: None.

TOMATO - For control of Colorado Potato Beetle, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetles. PHL: 7 days.

#### NOTICE OF WARRANTY

Kincaid Enterprises, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of K. E., Inc. In no case shall K. E., Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. K. E., INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.