

41014-3

PM-4

10-6-98

1/4



DRAFT

MARLATE GARDEN INSECTICIDE 5% DUST

For Effective Control Of Insects On Flowers, Fruits,
Vegetables, Ornamentals, Shade Trees, Shrubs,
And Dairy & Beef Cattle

Active Ingredient:

Methoxychlor, Technical*.....	5.0%
Other Ingredients.....	95.0%

*5.0% Methoxychlor Technical is equivalent to 4.75% 1,1,1-trichloro-2,2-bis(4-methoxyphenyl) ethane plus 0.25% impurities associated with the active ingredient.

EPA Registration No. 41014-3

EPA Establishment No. 41014-TN-001

KEEP OUT OF REACH OF CHILDREN

CAUTION


STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat.
 If on skin: Wash thoroughly with soap and water.
 If in eyes: Flush with plenty of water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET 1 or 4 LBS.

KINCAID ENTERPRISES, INC.
P.O. Box 549, Nitro, WV 25143



(back panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Do not apply this product through any type of irrigation system.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restrictive-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: Coveralls, Waterproof gloves, Shoes plus socks.

DIRECTIONS

Flowers, Fruits, Vegetables, Ornamentals, Shade Trees And Shrubs: Thoroughly dust the recommended amount of this product on foliage to cover all plant parts. Apply using conventional or other dusting equipment. Begin application at first sign of infestation (for fruits, begin at petal fall) and repeat at 7 - 14 day intervals or as needed to maintain control. Do not apply to crops within the number of days specified before harvest as shown by pre-harvest limitation (PHL).

PLANTS	INSECTS	Application Rate	PHL
Flowers & Ornamentals	Blister Beetles, Cankercworms, Cucumber Beetles, Flea Beetles, Fleahoppers, Flower Thrips, Japanese Beetle, Leafhoppers, Rose Chafer, Rose Slugs (Sawflies)	(1)	None
Shade trees and Shrubs	Tent Caterpillars, Cankercworms, Japanese Beetle	(1)	None
Apple, Pear, Quince	Apple Maggot, Codling Moth, Japanese Beetle, Plum Curculio, Tent Caterpillars	(1)	7 days
Apricot, Nectarine, Peach	Cherry Fruitworm, Cherry Fruit Flies, Japanese Beetle, Plum Curculio, Rose Chafer, Tent Caterpillars, Cankercworms	(1)	21 days
Cherry, Plum, Prune	Cherry Fruitworm, Cherry Fruit Flies, Japanese Beetle, Plum Curculio, Rose Chafer, Tent Caterpillars, Cankercworms	(1)	7 days
Blueberry, Currant, Gooseberry	Japanese Beetle, Cranberry Fruitworm, Leafhoppers, San Jose Scale (Crawlers), Strawberry Rootworm, Blueberry Flea Beetle Larvae, Blueberry Leaf Beetle, Blueberry Leaf Tier Larvae	(2)	14 days
Blackberry, Loganberry, Raspberry, Boysenberry, Dewberry, Youngberry, Strawberry	Rose Chafer, Strawberry Weevil, Flea Beetles, Omnivorous Leaf Tier, Spittlebugs, Japanese Beetle, Rednecked Cane Borer	(2)	14 days
Grape	Berry Moth, Grape Leafhopper, Japanese Beetle, Leaf Skeletonizer, Rose Chafer	(2)	14 days
Asparagus	Asparagus Beetle	(2)	(4)
Beans, Black-Eyed Peas	Corn Earworm, Cucumber Beetles, Fall Armyworm, Alfalfa Webworm, Garden Webworm, Flea Beetles, Mexican Bean Beetle, Potato Leafhopper, Bean Leaf Beetle, Japanese Beetle	(2)	(5)
Brussels Sprouts, Beets, Broccoli, Carrot, Collards, Kale, Lettuce, Radish, Spinach, Turnip	Blister Beetles, Flea Beetles, Leafhoppers, Alfalfa Looper, Fall Armyworm, Japanese Beetle, Imported Cabbageworm	(2)	14 days
Cauliflower, Eggplant, Kohlrabi, Pepper, Rutabaga	Blister Beetles, Flea Beetles, Leafhoppers, Alfalfa Looper, Fall Armyworm, Japanese Beetle, Imported Cabbageworm	(3)	7 days
Cabbage	Blister Beetles, Flea Beetles, Leafhoppers, Alfalfa Looper, Fall Armyworm, Japanese Beetle, Imported Cabbageworm	(2)	3 days
Sweet Corn	Armyworm, Fall Armyworm, Japanese Beetle, Flea Beetles	(2)	7 days
Cucumbers, Melons, Pumpkin, Squash, Summer Squash	Squash Vine Borer, Fall Armyworm, Cucumber Beetles, Flea Beetles	(2)	7 days
Peas	Alfalfa Looper, Pea Weevil (to foliage early bloom before eggs are laid and again 5 days later), Bean Leaf Beetle	(2)	7 days
Potato (Irish)	Colorado Potato Beetles, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetles	(1)	None
Sweet Potato, Yam	Fall Armyworm, Flea Beetles	(1)	None
Tomato	Colorado Potato Beetle, Flea Beetles, Fall Armyworm, Leafhoppers, Blister Beetles	(2)	7 days

Notes:

- (1) Dust on plant to obtain a full coverage application
- (2) Apply 4 ozs. per 100 square feet.
- (3) Apply 3 ozs. per 100 square feet.
- (4) If applied within 3 days of harvest, remove residues by washing or blanching.
- (5) 7 days; can be applied within 3 days of harvest if vines are not used for feed or forage.

DIRECTIONS (CONTINUED)

Dairy Cattle: Direct dusting for control of Hornfly. Sprinkle on poll, neck (top and sides) and back and upper sides; rub in lightly to carry powder beneath the hair. Repeat treatment every 3 weeks as needed. Do not use on dairy goats. Do not apply to lactating animals within five (5) hours of milking or during milking.

Beef Cattle, Sheep, Swine, Non-Lactating Dairy Cattle And Non-Lactating Goats: Direct dusting for control of Hornfly, Lice and Fleas. Dust animal thoroughly; rub in lightly to carry powder beneath the hair. Repeat treatment every 3 weeks as needed. May be used on young dairy animals up to two weeks of freshening. Application on sheep following shearing gives temporary relief from sheep ticks (keds).

Farm Buildings (except poultry houses): Direct dusting for control of stable flies, houseflies, fleas, gnats, mosquitoes, roaches and wasps. Apply dust to surface treating walls, floors, cracks, crevices, baseboards, pct bedding, stalls and pens. Such heavy deposits are usually effective for 3 to 6 weeks, unless removed by weathering or other means. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying. The use of any residual fly spray should be supplemented with proper sanitary measures.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

PRECAUTIONARY STATEMENTS: Caution - harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT: Handlers who may be exposed to this product through loading, application or other tasks must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Wash hands before, eating, drinking, chewing gum, using tobacco or using the toilet.

RED

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.


STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Reclose container after each use. Store in a cool, dry place away from food or feed. If spilled, sweep up and dispose of according to instructions shown below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container, then dispose of container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of the smoke.



(side panel)

4/4

NOTICE OF WARRANTY

Kincaid Enterprises, Inc. (hereinafter "Kincaid") warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Kincaid. In no case shall Kincaid be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks are assumed by the Buyer. KINCAID MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

