FEB 1 1994

Mr. Michael M. Watts Kincaid Enterprises, Inc. P.O. Box 549 Nitro, West Virginia 25143

Dear Mr. Watts:

Subject: KE Chloroneb 65W Fungicide

EPA Reg. No. 41014-4

Your Submission of November 3, 1993

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable. A stamped copy of the label is enclosed for your records.

Note that this labeling does not bear the required Worker Protection statements as outlined in PR Notices 93-7 and 93-11. We note that you have submitted labels in accordance with the instructions in those Notices. The new protective language in those submissions must be melded into this label for any shipments released after the deadlines that have been established. You may opt to do so immediately if you are electing the self-verification option, or after acceptance by the special Agency Worker Protection Standard review group. Finished printed labels should be submitted after acceptance by the WPS group and in accordance with their instructions.

Sincerely yours,

Sidney C. Jackson

Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

enclosure

7505C:C.Grable:cg:2/1/94

2405

ACCEPTED
with COMMENTS
In EPA Letter Dated

FEB 1 1994 Under the Federal Insecticide, Fundicide, and Redesticide Act as assessed, for the posticide registered under EPA Reg. No.



CHLORONEB 65W

FUNGICIDE

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Flush with plenty of water.

SEE THE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

GENERAL INFORMATION—K.E. "Chloroneb" 65W Fungicide is recommended for use as a supplemental seed treatment, for the control of seedling diseases of cotton, beans and soybeans; and as a supplemental seed treatment for sugar beets. It is to be used in conjunction with a standard fungicide seed treatment.

When used as directed, "Chloroneb" 65W can be expected to improve stands and plant vigor through the suppression of seedling blights, soreshin, and pre- and post-emergence damp-off caused by such organisms as Rhizoctonia and Pythium species and Sclerotium rolfsii. It is highly effective in preventing post-emergence damp-off, especially when used as a supplemental seed coating.

BEST AVAILABLE COPY

Net Weight: 25 Lbs

Mitg.
KINCAID ENTERPRISES, INC.

BOX 549 NITRO, WV 25143

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

3015

PRECAUTIONARY STATEMENTS: Caution—harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT (PTE): Handlers who may be exposed to the dilute through application of other tasks must wear:

Long-sleeved shirt and long pants

Waterproof gloves

Shoes plus socks

Handlers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear:

Long-sleeved shirt and long pants

Waterproof gloves

Shoes plus socks

USER SAFETY RECOMMENDATIONS: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Reclose container after each use. Store in a cool, dry place away from food or feed. If spilled sweep up and dispose of below.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag, then dispose of bag in a sanitary landfill or by inceneration, or, if allowed by state and local authorities, by burning. If burned, stay out of the smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on tarms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for trining, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restrictive-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated a eas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls

Waterproof gloves

Shoes plus socks

BEST AVAILABLE COPY

Stocked.

Do not apply this product through any type of irrigation system.

SLURRY TREATERS

CHLORONEB 65W may be applied separately as a seed overcoat or in combination with other standard seed protectants. Mix CHLORONEB 65W in water and apply at the rates, slurry cup sizes and dump weights shown below. When using CHLORONEB 65W in combination with other seed protectants, adjust the slurry cup sizes and dump weights for the increased amount of water needed to handle the additional solids. Observe all cautions and limitations on labeling of all products used in mixtures. The treating rates for cottonseed include acid delinted, reginned, or fuzzy cottonseed. Use 4 pounds CHLORONEB 65W per gallon of water; one fluid ounce equals approximately 29.5 cc.

Crop Seed	Treating Rate Oz/CWT	Slurry cc/CWT Seed	Slurry Cup, cc	Lbs Dump Weight
Cotton - East of Rock	cies 6	460	46	10
Cotton - West of Rock	cies 10	766	46	6
Cotton (High Plains area of '	4 Texas)	306	46	6.7
Beans and Soybeans	4	306	23	7.5
Sugar Beets	6	460	46	10

HOPPER BOX TREATMENT

The following procedure is suggested to thoroughly mix CHLORONEB 65W with the seed before planting: premix one-half the required solution at the rate of 4 lbs CHLORONEB 65W per gallon of water and fill planter box half full with seed. Add the premixed solution of CHLORONEB 65W and mix thoroughly with a stick. Prepare the second half of the CHLORONEB 65W as described above. Then add the remainder of the seed to the planter box and add the rest of the required amount of CHLORONEB 65W to the seed, again mixing thoroughly.

Crop Seed	Treating Rate Oz/CWT	Slurry cc/CWT Seed	
Cotton (East of Rockies)	6	460	•
Cotton (West of Rockies)	10	766	
Cotton (High Plains - TX)	4	306	' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '
Beans and Soybeans	4	306	•
Sugar Beets	6	460	

NOTICE: All stored treated seed must be labeled: "This seed treated with Chloroneb 65W fungicide. Do not use for food, feed, or oil purposes." Add appropriate information for any other chemical treatments. Do not use excess treated seed for food, feed, or oil purposes. Do not graze treated beans or soybeans within 45 days of planting.