





ACTIVE INGREDIENT:	
Chloroneb (1,4-dichloro-2,5-dimethoxybenzene).	65%
INERT INGREDIENTS	
U.S. Pat. 3,265,534	EPA Reg. No. 41014-6

GENERAL INFORMATION—Terraneb SP Turf Fungicide is recommended for the treatment of turfgrasses for the control of snow mold (Typhula) and Pythium blight.

Keep out of reach of children.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! MAY IRRITATE EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. First Aid: In case of contact, flush skin or eyes with plenty of water: for

# eyes, get medical attention

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated

# PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire or sparks.

# REREGISTERED

ACCEPTED with COMMENTS in EPA Litter Untest:

SEP 2 9 1982

361 2 3 1301
Under the content I.
Funge to a cities
ra bull dia, and
rg 1354 # 1 1.44 1
4/014-6

Made in U.S.A

NET 3 LBS.

EPA Est. 42761-MS 1



### -NOTICE OF WARRANTY

K. E. Inc. warrants that this product centerms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of K. E. Inc. In no case shall K. E. Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer K. E. INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

## **DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### TURF DISEASES

Snow Mold—For the control of snow mold (Typhula), apply 6 to 9 ozs. of Terraneb SP per 1000 sq. ft. in sufficient water to give complete coverage, usually 3 to 5 gallons. Apply as a spray as late in the fall as possible prior to snowfall.

Pythlum Blight—For control of Pythium blight on established turf, apply Terraneb SP as a preventative at a rate of 4 ozs. per 1000 sq. ft. in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of disease. Repeat in 5 to 7 days if conditions persist.

For the control of Pythium blight on new seedings, apply Terraneb SP at the rate of 4 ozs. per 1000 sq. ft. following emergence. If conditions remain favorable to the development of disease, repeat in 7 to 10 days.

Note to User: Do not graze or feed clippings from treated areas to livestock.

NOTICE TO BUYER-Purchase of this material does not confer any rights under patents of countries outside of the United States.

# STORAGE AND DISPOSAL

PROHIBITIONS — Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Avoid contamination of feed and food-stuff.

PESTICIDE DISPOSAL— Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures

CONTAINER DISPOSAL — Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner and drum in a sanitary landfill or by incineration if allowed by state and local authorities.