



199  
PM 21  
41014-6  
Pg 1 of 3

# TERRANE



# SP

## TURF FUNGICIDE

**ACTIVE INGREDIENT:**

Chloroneb (1, 4-dichloro-2, 5-dimethoxybenzene) ..... 65%

**INERT INGREDIENTS** ..... 35%

U.S. Pat. 3,265,564

EPA Reg. No. 41014-6

EPA Est No. 35982-TX-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Call a Physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash the affected area with soap and warm water.

**IF IN EYES:** Flush with plenty of water. Call a Physician.

See side panel for additional precautionary statements.

**ACCEPTED**

**AUG 28 1987**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 41014-6

## NET 3 LBS.

**Caution:** Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**Storage:** Reclose container after each use. Store in a cool, dry place away from food or feed. If spilled, sweep up and dispose of below.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty can, then dispose of can in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### NOTICE OF WARRANTY

Kincaid Enterprises, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on this label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Kincaid Enterprises, Inc. In no case shall Kincaid Enterprises, Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Kincaid Enterprises, Inc. makes no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



**KINCAID ENTERPRISES**

P. O. BOX 671 • NITRO, WV 25143

## TURFGRASS DISEASES

Typhula Blight (Gray Snow Mold) - For control of Typhula Blight, apply 6 to 9 ozs. TERRANE SP per 1000 sq. ft. in sufficient water to give complete coverage, usually 3 to 5 gallons. Apply as a spray as late in the fall as possible prior to snowfall.

Pythium Blight (Cottony Blight) and Sclerotium Blight - For control of Pythium Blight and Sclerotium Blight on established turf, apply TERRANE SP as a preventative at a rate of 4 ozs. per 1000 sq. ft. in 3 to 5 gallons of water. Apply during periods of high temperature and humidity favorable to the development of the diseases. Repeat in 5 to 7 days if conditions persist.

For control of Pythium Blight and Sclerotium Blight on new seedlings, apply TERRANE SP at the rate of 4 ozs. per 1000 sq. ft. following emergence. If conditions remain favorable to the development of the diseases, repeat in 7 to 10 days.

Rhizoctonia Blight (Brown Patch) - For warm season lawns and turfs (such as zoysiagrass, St. Augustine, bermudagrass and similar grasses) apply in the fall or spring at the first signs of infection. Mix 5 ozs. TERRANE SP in 10 - 15 gallons water to thoroughly cover 1000 sq. ft. Make a second application in 3 - 4 weeks if the disease is severe or re-appears.

For cool season lawns and turf (such as creeping bentgrass, annual bluegrass, fescue and similar grasses) apply at first signs of infection. Mix 3 - 4 ozs. TERRANE SP in 3 - 5 gallons water to thoroughly cover 1000 sq. ft. Reapply every week to 10 days during warm, damp weather.

### NOTICE

Do not graze or feed clippings from treated areas to livestock.

Mfg:

KINCAID ENTERPRISES, INC.  
Box 671, Nitro, WV 25143