

41052-9
ACCEPTED
 JUL 8 1960
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended for the purpose of registration under EPA Reg. No. 41052-9

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations

For sanitizing food processing equipment, dairy equipment, food service dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. No Potable water flow is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. CCC Disinfectant-Sanitizer-Fungicide-Deodorizer in 4 gallons of water.

Apply to sink tops, counters, refrigerated storage and display equipment and other necessary and hard surfaces by cloth or brush. No Potable water flow is required.

Dishes, silverware, glasses, cooking utensils and other similar food processing equipment can be sanitized by immersion in a 1 oz. 4 gallon solution of CCC Disinfectant-Sanitizer-Fungicide-Deodorizer. No Potable water flow is required.

At 1 oz. 4 gallons, CCC Disinfectant-Sanitizer-Fungicide-Deodorizer fulfills the criteria of Appendix C of the Grade A Pasteurized Milk Ordinance 1965 Recommendations, one of the U.S. Public Health Service in waters up to 1000 ppm of hardness, calculated as Ca CO₃ when evaluated by U.S. AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, teats and teats of dairy cows can be sanitized by washing with a solution of 1 oz. CCC Disinfectant-Sanitizer-Fungicide-Deodorizer in 4 gallons of warm water. No Potable water flow is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements Hazards to Humans and Domestic Animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.
 - OPEN DUMPING IS PROHIBITED.
 - DO NOT REUSE EMPTY CONTAINER.

HAZARDOUS DISPOSAL
 PESTICIDE OR HERBICIDE THAT CANNOT BE USED OR CHEMICALLY REPRODUCED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURNED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL
 TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

GENERAL
 CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

CCC DISINFECTANT-SANITIZER FUNGICIDE DEODORIZER

Disinfectant-Sanitizer-Fungicide Deodorizer for Hospital, Institutional, Industrial, School, Dairy and Other Farm and Home Use

AOAC Phenol Coefficients

Staphylococcus aureus, ATCC #6516 77.5
Salmonella typhosa, ATCC #6539 77.5

Active Ingredients

Dicetyl dimethyl ammonium chloride 7.5
 Isopropanol 30

Inert Ingredients

89.5
 100.00%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 41052-9
 Net Contents

Est No 41052-Me-1

Manufactured By
 Chute Chemical Company, Bldg 97, B I A, Bangor, Maine 04401

Directions for Use

GENERAL CLASSIFICATION

In accordance with Federal Law to use this product in a manner consistent with its labeling.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 1 oz. CCC Disinfectant-Sanitizer-Fungicide-Deodorizer to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use level, CCC Disinfectant-Sanitizer-Fungicide-Deodorizer is effective against *Pseudomonas aeruginosa*.

Disinfection in Institutions, Industry, Schools and Homes

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 2 oz. of CCC Disinfectant-Sanitizer-Fungicide-Deodorizer to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz. 4 gallon use level, CCC Disinfectant-Sanitizer-Fungicide-Deodorizer is effective against *Staphylococcus aureus*, *Escherichia coli*, *Salmonella choleraesuis* and *Trichophyton mentagrophytes* (the athlete's foot fungus).

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1-oz. 4 gallon solution of CCC Disinfectant-Sanitizer-Fungicide-Deodorizer.

Disinfection of Poultry Equipment, Animal Quarters and Kennels

Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. CCC Disinfectant-Sanitizer-Fungicide-Deodorizer in 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.