

ACCEPTED
 DEC - 2 1985
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. *4124-5*



25-5-10 WITH Team™

Active Ingredient	
N-butyl-N-ethyl- α,α,α -trifluoro-2,6-dinitro-p-toluidine	74%
trifluralin: α,α,α -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	3%
Inert Ingredients	98.87%

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impervious gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention.

Environmental Hazards

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Storage: Store in original container only. In case of leak or spill, contain material and dispose of waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

E.P.A. REG. NO. 41124-

E.P.A. EST. NO. 41124-WI-1

NET WT. 50#

COVERS 12,100 SQ. FT.

MFG. BY

Vogel Seed & Fertilizer, Inc.
Jackson, WI 53037

SPRING VALLEY'S 25-5-10 with TEAM is a high quality fertilizer product which combines turf fertilizer, a pre-emergence herbicide and a selective pre-emergence herbicide to control the below listed annual grasses.

DIRECTIONS FOR USE Read all Directions Carefully Before Applying

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. SPRING VALLEY'S 25-5-10 with TEAM is a pre-emergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (spoon annual), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), banyardgrass (watergrass) and green and yellow foxtail. SPRING VALLEY'S 25-5-10 with TEAM controls these annual weeds by killing their seeds as they germinate. Apply before these annual weeds begin to germinate since it does not control established weeds.

When SPRING VALLEY'S 25-5-10 with TEAM is applied as directed, the following established turfgrasses are tolerant:

Kentucky bluegrass	Fescue	St. Augustinegrass
Perennial ryegrass	Zoysiagrass	Bahiagrass
Centipede grass	Bermudagrass	

SPRING VALLEY'S 25-5-10 with TEAM will not harm most nearby established ornamental trees and shrubs.

APPLICATION DIRECTIONS

Rates, Frequency and Timing of Application

Successful pre-emergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply SPRING VALLEY'S 25-5-10 with TEAM approximately one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance will be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application of SPRING VALLEY'S 25-5-10 with TEAM.

Spring germination summer annual grasses-crabgrasses, foxtails and banyardgrass

One 50# bag of SPRING VALLEY'S 25-5-10 with TEAM will cover 12,100 sq. ft. or 180# per acre. Application at this rate will supply 1# of actual nitrogen per 1000 sq. ft. and 2# of actual TEAM per acre. Do not reseed 12 to 16 weeks after application.

A. Northern, cool season turfgrasses
For pre-emergence control, apply 175 to 190 pounds per acre (1.5 to 2 pounds active ingredient) of SPRING VALLEY'S 25-5-10 with TEAM per acre in the spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

B. Southern, warm season turfgrasses
For pre-emergence control, apply 190 to 270 pounds (2 to 3 pounds active ingredients) of SPRING VALLEY'S 25-5-10 with TEAM per acre in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of SPRING VALLEY'S 25-5-10 with TEAM at the recommended rate may be made 8 to 10 weeks after initial treatment for continued control.

SPRING VALLEY'S 25-5-10 with TEAM Per Acre	#/ 1000 Sq. Ft.
175 pounds	3.1 pounds
190 pounds	4.1 pounds
270 pounds	6.2 pounds

PRECAUTIONS

Apply SPRING VALLEY'S 25-5-10 with TEAM to the turf prior to the germination of weed grasses since SPRING VALLEY'S 25-5-10 with TEAM will not control established weeds. SPRING VALLEY'S 25-5-10 with TEAM may thin established annual bluegrass (spoon annual) turfs and fine leaved fescues at rates above 125 pounds per acre (1.5 pounds active ingredient). SPRING VALLEY'S 25-5-10 with TEAM should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of SPRING VALLEY'S 25-5-10 with TEAM may thin the overseeded grasses. Because of varying cultural practices, SPRING VALLEY'S 25-5-10 with TEAM should not be used on putting greens. SPRING VALLEY'S 25-5-10 with TEAM should not be used on diondrea. SPRING VALLEY'S 25-5-10 with TEAM should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass or Centipede grass until these turfs are well established.

GUARANTEED ANALYSIS

Nitrogen (N).....25%
 10.5% urea nitrogen
 2.0% ammoniacal nitrogen
 12.5% sulfur coated urea (SCU)
 Available Phosphorus (acid TP25).....5%
 Soluble Potash (K2O).....10%
 Sulfur.....5%

FERTILIZER INGREDIENTS: sulfur coated urea, urea, D.A.P., muriate of potash.

NET WT. 50# COVERS 12,100 SQ. FT.

SPRING VALLEY TURF PRODUCTS
 1841 Spring Valley Rd. Jackson, WI 53037
 Div. Vogel Seed & Fert.